

Poster presentation

Open Access

An increased N-methyl-D-aspartate receptor conductance is associated with intrinsic bursting behavior

Amber Martell*¹, Seoan Marler¹, Hyong C Lee¹, Jan-Marino Ramirez² and Wim van Drongelen¹

Address: ¹Department of Pediatrics, University of Chicago Hospitals, The University of Chicago, Chicago, IL, USA and ²Department of Organismal Biology and Anatomy, The University of Chicago, Chicago, IL, USA

Email: Amber Martell* - martella@uchicago.edu

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P144 doi:10.1186/1471-2202-8-S2-P144

© 2007 Martell et al; licensee BioMed Central Ltd.

Background

Seizure activity is often accompanied by an increase in the number of intrinsically bursting neurons [1]. The N-methyl-D-aspartate (NMDA) receptor, a channel known to be involved in many seizure models, represents one mechanism by which intrinsic bursting may arise, since 30–50% of these channels are bound at ambient concentrations of glutamate. In adult rat, intrinsic bursting has been induced in pyramidal neurons of the prefrontal cortex (PFC) through a combination of NMDA and dopamine type 1 (D1) receptor agonist [2]. Here we used the in vitro mouse neocortex and a computational model to further investigate the basis for bursting in the context of epileptiform activity.

Results

In mouse frontal cortex, application of 8–10 μM NMDA and 2–5 μM D1 agonist SKF 38393 elicited intrinsic bursting in $\sim 50\%$ of pyramidal neurons. A computational pyramidal neuron model consisting of five compartments with the sodium, potassium, calcium-activated potassium and NMDA channels was used to investigate the conditions necessary for bursting. In this model D1 agonist caused an amplification of the NMDA current. Our simulations indicate that increasing the NMDA receptor conductance transformed a regularly spiking neuron into a burster (see fig. 1). Interestingly, the bursting behavior




Figure 1
Experimental and Model Neuron Firing Behaviour.

appeared only when both NMDA receptor and calcium-activated potassium conductance were included.

Conclusion

The NMDA receptor is associated with production of intrinsic bursting behavior in mouse cortical pyramidal neurons, and, together with a calcium-activated K conductance, is sufficient to cause spontaneous bursting in a computational model of pyramidal neuron. These and previous results indicate that the NMDA receptor has the potential to drive the bursting behavior that characterizes seizures.

Acknowledgements

This work was supported in part by the Falk Foundation and the Linn family.

References

1. Topolnik L, Steriade M, Timofeev I: **Hyperexcitability of intact neurons underlies acute development of trauma-related electrographic seizures in cats in vivo.** *Eur J Neurosci* 2003, **18**:486-96.
2. Durstewitz D, Gabriel T: **Dynamical Basis of Irregular Spiking in NMDA-Driven Prefrontal Cortex Neurons.** *Cerebral Cortex* 2006 in press.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

