

Poster presentation

Open Access

Dopamine D1/D2 modulation of synaptic plasticity in the prefrontal cortex

Robert Kyle*¹, Ullrich Bartsch², David Willshaw¹ and Daniel Durstewitz³

Address: ¹Institute for Adaptive and Neural Computation, University of Edinburgh, Edinburgh, EH8 9AB, UK, ²Department of Physiology and Pharmacology, University of Bristol, Bristol, BS8 1TD, UK and ³Central Institute for Mental Health Mannheim, University of Heidelberg, Mannheim, Germany

Email: Robert Kyle* - r.kyle@ed.ac.uk

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, 10(Suppl 1):P193 doi:10.1186/1471-2202-10-S1-P193

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P193>

© 2009 Kyle et al; licensee BioMed Central Ltd.

Introduction

It is widely assumed that dopamine plays a crucial role in reinforcement learning due to the ability of dopamine neurons to signal reward prediction errors. However, it is not clear how such a signal could affect neurons in the prefrontal cortex and whether or not the experimentally demonstrated effects of dopamine on long-term synaptic plasticity are by themselves enough to cause the animal to learn to predict future rewards. The aim of this study was to investigate the effect of dopamine modulation on synaptic plasticity via both the D1 and D2 dopamine receptors.

Methods

In this study, the effect of dopamine modulation on synaptic plasticity was determined by simulating a selection of plasticity protocols under both control conditions and a simulated bath of D1 or D2 agonists. The effects of dopamine D1 and D2 agonists on both intrinsic and synaptic parameters of PFC neurons were quantified using data obtained from in vitro recordings. In addition to the effects of dopamine on the single neuron *f/I* curve, D1 agonists enhanced calcium influx, NMDA and GABAA currents in the model whilst reducing the effect of AMPA conductances. D2 agonists acted oppositely on calcium influx and synaptic conductances. Changes in synaptic efficacy were quantified using an existing calcium-based model of synaptic plasticity [1].

Results

In states where D1 modulation dominated, there was enhanced LTP of active synapses, an effect that is attenuated by D2 receptor stimulation, which by itself results in less potentiation, or an increased likelihood of LTD. The effect of dopamine modulation on STDP showed a complex timing-dependence and cannot be simply interpreted in terms of an increase or decrease in plasticity (see Figure). Together our results indicate that calcium dependent models of plasticity are highly sensitive to neuromodulators which effect calcium influx, and therefore a more detailed understanding of single neuron calcium dynamics will be required to accurately characterize the effect of dopamine on synaptic plasticity.

References

1. Shouval HZ, Bear MF, Cooper LN: **A unified model of NMDA receptor-dependent bidirectional synaptic plasticity.** *Proc Natl Acad Sci USA* 2002, **99**:10831-10836.




Figure 1
The effect of dopamine modulation on STDP in a frequency dependent plasticity protocol.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

