

Poster presentation

Open Access

Partial synchronization in diluted neural networks

Simona Olmi*^{1,3}, Roberto Livi^{1,3}, Antonio Politi² and Alessandro Torcini^{2,3}

Address: ¹Department of Physics, University of Florence, Sesto Fiorentino, I-50019, Italy, ²Istituto dei Sistemi Complessi, CNR, Sesto Fiorentino, I-50019, Italy and ³Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, Sesto Fiorentino, I-50019, Italy

Email: Simona Olmi* - simona.olmi@fi.isc.cnr.it

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P274 doi:10.1186/1471-2202-10-S1-P274

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P274>

© 2009 Olmi et al; licensee BioMed Central Ltd.

Partial synchronization (PS) is a collective dynamics discovered in fully coupled networks of leaky integrate-and-fire (LIF) neurons [1,2]. This dynamical state is quite peculiar, since the mean-field is characterized by a periodic dynamics, while the single neurons do not show any synchronization and behaves quasi-periodically in time. PS arises from the destabilization of a regime characterized by a constant mean field and by periodic motion of the single neurons. This regime, termed "Splay State," has been widely studied in several contexts, including computational neuroscience [3].

Our aim is to investigate the role of network dilution on the stability properties of PS and splay states and the influence of dilution on the transition scenario leading from the splay state to partial synchronization. In particular, we have examined a pulse-coupled excitatory network of LIF neurons with 20% of randomly broken links in two different configurations: quenched, where the network topology is fixed, and annealed, where the links to break are chosen at each spike emission. For both configurations, we have been able to rewrite the dynamics of the network (described by a set of differential equations) as an exact event driven map by extending the approach developed in [4].

We observe that the splay state and the PS state, as well as the transition scenario, are quite robust with respect to dilution, both for the quenched and annealed configurations. Moreover the dynamics for finite number of neurons is chaotic at variance with the fully coupled case. The

dependence of the degree of chaotic behavior (characterized in terms of the maximum Lyapunov exponent) on the number of neurons N is analyzed, as a first result we notice that finite- N effects are more relevant in the quenched configuration with respect to the annealed one.

Finally, the possibility to reproduce the deterministic dynamics of the diluted network for finite N , by introducing an additive colored noise term in the equations describing the fully coupled case, is investigated. The noise term amplitude and correlation are determined by the analysis of the mean field properties associated with finite diluted networks.

References

1. van Vreeswijk C: **Partial synchronization in populations of pulse-coupled oscillators.** *Phys Rev E* 1996, **54**:5522-5537.
2. Mohanty PK, Politi A: **A new approach to partial synchronization in globally coupled rotators.** *J Phys A* 2006, **39**:L415-L420.
3. Abbott LF, van Vreeswijk C: **Asynchronous states in networks of pulse-coupled oscillators.** *Phys Rev E* 1993, **48**:1483-1490.
4. Zillmer R, Livi R, Politi A, Torcini A: **Stability of the splay state in pulse-coupled networks.** *Phys Rev E* 2007, **76**:046102.