

POSTER PRESENTATION

Open Access

The impact of pooling and shared inputs on correlations in neuronal networks

James Trousdale*, Robert Rosenbaum, Krešimir Josíc

From Nineteenth Annual Computational Neuroscience Meeting: CNS*2010 San Antonio, TX, USA. 24-30 July 2010


A cortical neuron receives inputs from thousands of afferents. Experiments suggest that the activity of these afferent cells is often correlated, although such correlations may be weak. Similarly, recordings of field potentials or data obtained using voltage sensitive dyes represents the pooled activity of large populations of cells, which can be correlated. It is therefore important to understand how correlations between cells in a population affect the statistics of the pooled activity of cells taken from the population.

It is well known that even weak correlations within an input pool can increase the variability of the pooled signal. This phenomenon and its implications for the response and coding capabilities of a downstream cell were investigated in [1][2][3].

Here we address a related phenomenon: weak correlations within and between *two* populations of neurons

lead to significant correlations between the two pooled signals [4]. This phenomenon has been observed in the context of correlations between two VSD [5] and MUA [6] signals. We focus primarily on the effect pooling in input populations has on the covariation of the membrane potentials of two downstream cells. We use simple probabilistic formulae to provide an intuitive explanation for this phenomenon, and show that the contribution of overlap in input populations to the development of correlations is minor relative to the contribution of pooling.

In addition, we apply our results to provide insight on development of correlations and synchrony in feed-forward networks. We show that pooling, and not overlap in input populations, contributes most to the development of very high correlations observed in feed-forward networks. We exhibit via simulations that not only cor-



^{*} Correspondence: jrtrousd@math.uh.edu Department of Mathematics, University of Houston, Houston, TX 77004, USA



relations, but also synchrony, develop in the absence of overlap in the input populations. (Figure 1) In a related example, we show that in more structured feed-forward networks including local disynpatic inhibitory circuits, moderate synchrony (i.e., correlations over short time windows) is a strong requirement for the propagation of inputs.

Published: 20 July 2010

References

- Shadlen MN, Newsome WT: The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. J Neurosci 1998, 18:3870-3896.
- Salinas E, Sejnowski TJ: Impact of correlated synaptic input on output firing rate and variability in simple neuronal models. J Neurosci 2000, 20:6193-6209.
- Moreno-Bote R, Renart A, Parga N: Theory of input spike auto- and crosscorrelations and their effect on the response of spiking neurons. Neural Comput 2008. 20:1651-1705.
- Renart A, de la Rocha J, Bartho P, Hollender L, Parga N, Reyes A, Harris KD: The asynchronous state in cortical circuits. Science 2010, 327:587-590.
- Chen Y, Geisler WS, Seidemann E: Optimal decoding of correlated neural population responses in the primate visual cortex. Nat Neurosci 2006, 9:1412-1420.
- Stark E, Globerson A, Asher I, Abeles M: Correlations between groups of premotor neurons carry information about prehension. J Neurosci 28:10618-10630.

doi:10.1186/1471-2202-11-S1-P12

Cite this article as: Trousdale *et al.*: The impact of pooling and shared inputs on correlations in neuronal networks. *BMC Neuroscience* 2010 11 (Suppl 1):P12.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

