

ORAL PRESENTATION

Open Access

# A cell-type-specific dynamic Bayesian network model for spontaneous and optogenetically evoked activity in the primary visual cortex

Ali Mohebi<sup>1</sup>, Jessica A Cardin<sup>3</sup>, Karim G Oweiss<sup>1,2\*</sup>

From Twentieth Annual Computational Neuroscience Meeting: CNS\*2011  
Stockholm, Sweden. 23-28 July 2011

Reciprocal interaction between excitatory and inhibitory neurons within and between layers of the cerebral cortex is a major element of brain function. Ensemble extracellular recording techniques using microelectrode arrays have permitted recording spiking activity of many neurons simultaneously to characterize network function [1]. Identifying the type of neurons in these recordings is not straightforward due to the variability in extracellular spike shapes, and the irregularities often observed in their interspike interval characteristics. In this study, we used optogenetic tools to genetically target fast spiking interneurons in the primary visual cortex of mice [2]. We modulated their spiking activity by illuminating the region with very short pulses (<1 ms) of light (~470 nm wavelength) in mice primary visual cortex. Using Dynamic Bayesian Network analysis [3], we assessed the effective connectivity between the recorded neurons in the presence and absence of light stimuli under distinct cortical states observed under light anesthesia. DBNs could identify the effective connectivity between putative excitatory pyramidal cells and inhibitory interneurons. These findings suggest a novel and unprecedented way to identify cortical neuronal circuits and characterize the dynamics of their computations *in vivo*.

## Author details

<sup>1</sup>Department of Electrical and Computer Engineering, Michigan State University, East Lansing, MI 48824, USA. <sup>2</sup>Neuroscience Program, Michigan State University, East Lansing, MI 48824, USA. <sup>3</sup>Department of Neurobiology, Yale University School of Medicine, New Haven, CT 06510, USA.

Published: 18 July 2011

\* Correspondence: koweiss@msu.edu

<sup>1</sup>Department of Electrical and Computer Engineering, Michigan State University, East Lansing, MI 48824, USA

Full list of author information is available at the end of the article

## References

1. Haider B, Duque A, Hasenstaub AR, McCormick DA: **Neocortical Network Activity In Vivo Is Generated through a Dynamic Balance of Excitation and Inhibition.** *Journal of Neuroscience* 2006, **26**:4535-4545.
2. Cardin JA, Carlen M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai L, et al: **Targeted optogenetic stimulation and recording of neurons in vivo using cell-type-specific expression of Channelrhodopsin-2.** *Nature Protocols* 2010, **5**:247-254.
3. Eldawlaty S, Zhou Y, Jin R, Oweiss KG: **On the Use of Dynamic Bayesian Networks in Reconstructing Functional Neuronal Networks from Spike Train Ensembles.** *Neural Computation* 2010, **22**:158-189.

doi:10.1186/1471-2202-12-S1-O5

**Cite this article as:** Mohebi et al: A cell-type-specific dynamic Bayesian network model for spontaneous and optogenetically evoked activity in the primary visual cortex. *BMC Neuroscience* 2011 **12**(Suppl 1):O5.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

