

POSTER PRESENTATION

Open Access

Synchronization hubs may arise from strong rhythmic inhibition during gamma oscillations in primary visual cortex

Stefanos E Folias^{1*}, Danko Nikolić^{2,3}, Jonathan E Rubin¹

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Parallel multiunit recordings from V1 in anesthetized cat were collected during the presentation of random sequences of drifting sinusoidal gratings at 12 fixed orientations while gamma oscillations were present. In agreement with the seminal work [1], most units were orientation selective to varying degrees and synchronization was evident in spike train crosscorrelograms computed between units with similar preferred orientations, particularly during the presentation of optimal stimuli. Interestingly, a subset of units, which we refer to as synchronization hubs, were additionally found to synchronize with units having differing preferred orientations which was consistent with a previous study [2]. Moreover, oscillatory patterning in spike train autocorrelograms was also found to be strongest in units denoted as synchronization hubs, and synchronization hubs also tended to have narrower tuning curves relative to other units.

We used simplified computational models of small networks of V1 neurons to demonstrate that neurons subject to a sufficiently strong level of inhibitory input can function as synchronization hubs. Neurons were endowed either with integrate-and-fire or conductance-based dynamics and each neuron received a combination of excitatory (AMPA) synaptic inputs that were Poisson-distributed and inhibitory (GABA) inputs that were coherent at a gamma-frequency range. If the strength of rhythmic inhibition was increased for a subset of neurons in the network, and excitation was increased simultaneously to maintain a fixed firing rate, then these neurons produced stronger oscillatory patterning in their discharge probabilities. The oscillations in turn synchronized these neurons with other neurons

in the network. Importantly, the strength of synchronization increased with neurons of differing orientation preferences even though no direct synaptic coupling existed between the hubs and the other neurons.

Enhanced levels of inhibition account for the emergence of synchronization hubs in the following way: Inhibitory inputs exhibiting a gamma rhythm determine a time window within which a cell is likely to discharge. Increased levels of inhibition narrow down this window further simultaneously leading to (i) even stronger oscillatory patterning of the neuron's activity and (ii) enhanced synchronization with other neurons. This enables synchronization even between cells with differing orientation preferences. Additionally, the same increased levels of inhibition may be responsible for the narrow tuning curves of hub neurons. In conclusion, synchronization hubs may be the cells that interact most strongly with the network of inhibitory interneurons during gamma oscillations in primary visual cortex.

Author details

¹Department of Mathematics, University of Pittsburgh, Pittsburgh, PA, USA.

²Department of Neurophysiology, Max Planck Institute for Brain Research, Wolfgang Goethe University, 60528 Frankfurt/Main, Germany. ³Frankfurt Institute for Advanced Studies, Wolfgang Goethe University, 60438 Frankfurt/Main, Germany.

Published: 18 July 2011

References

1. Gray C, Singer W: Stimulus-specific neuronal oscillations in orientation columns of cat visual cortex. *Proc. Natl. Acad. Sci* 1989, **86**:1698-1702.
2. Yu S, Huang D, Singer W, Nikolić D: A small world of neuronal synchrony. *Cerebral Cortex*. 2008, **18**:2891-2901.

doi:10.1186/1471-2202-12-S1-P277

Cite this article as: Folias et al.: Synchronization hubs may arise from strong rhythmic inhibition during gamma oscillations in primary visual cortex. *BMC Neuroscience* 2011 **12**(Suppl 1):P277.

* Correspondence: stefanos@pitt.edu

¹Department of Mathematics, University of Pittsburgh, Pittsburgh, PA, USA
Full list of author information is available at the end of the article