

POSTER PRESENTATION

Open Access

Integration of predictive-corrective incompressible SPH and Hodgkin-Huxley based models in the OpenWorm in silico model of *C. elegans*

Michael Vella^{1*}, Andrey Palyanov², Padraig Gleeson³, Sergey Khayrulin²

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

OpenWorm is an international collaboration with the aim of producing an integrative computational model of *Caenorhabditis elegans* to further the understanding of how macroscopic behaviour of the organism emerges from aggregated biophysical processes. A core component of the project involves the integration of electrophysiological modelling and predictive-corrective incompressible smoothed particle hydrodynamics (PCISPH) to model how neuronal and muscle dynamics effect the nematode's behaviour. Several tools are being utilised and developed in the course of the project:

- Electrophysiological model parameters are constrained to reproduce experimental measurements using the Optimal Neuron toolkit [1]
- A PCISPH solver is under development [2] - a combination of general PCISPH algorithms proposed by [3], boundary-handling algorithms proposed by [4], a surface tension model based on [5] and our own implementation of elastic matter and biophysics-specific features, as well as parallelization, optimization and tuning. It is the first open source, parallel OpenCL/C++ PCISPH high-performance implementation.
- A generic model integration framework (Gepetto [6]) will be used to integrate electrophysiology and body-wall interactions
- All electrophysiological models are NeuroML-compatible [7].

- The Open Worm Browser provides a powerful way to visualise *C. Elegans* anatomy [8]

All of the above mentioned applications are open source, freely available and can be used for modelling other neuronal systems.

Author details

¹Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, CB2 3DY, UK. ²Lab. of complex systems simulations, A.P. Ershov Institute of Informatics Systems, Siberian Branch of the Russian Academy of Sciences, Novosibirsk, 630090, Russian Federation. ³Department of Neuroscience, Physiology and Pharmacology, University College London, London, UK.

Published: 8 July 2013

References

1. Optimal Neuron Toolkit. [https://github.com/vellamike/Optimal-Neuron].
2. Open Worm PCISPH Simulator. [https://github.com/openworm/Smoothed-Particle-Hydrodynamics].
3. Solenthaler B, Pajarola R: Predictive-corrective incompressible SPH. *ACM Trans Graph* 2009, **28**(3).
4. Ihmsen M, Akinci N, Gissler M, Teschner M: Boundary Handling and Adaptive Time-stepping for PCISPH. *Proc VRIPHYS* Copenhagen, Denmark; 2010, 79-88, Nov 11-12.
5. Becker M, Teschner M: Weakly compressible SPH for free surface flows. *Proceedings of the 2007 ACM SIG-GRAPH/Eurographics* 2007.
6. Gepetto Simulation Engine. [https://github.com/openworm/OpenWorm/wiki/Gepetto-Overview].
7. Gleeson P, Crook S, Cannon RC, Hines ML, Billings GO, et al: NeuroML: A Language for Describing Data Driven Models of Neurons and Networks with a High Degree of Biological Detail. *PLoS Comput Biol* 2010, **6**(6): e1000815, doi:10.1371/journal.pcbi.1000815.
8. Open Worm Browser. [http://browser.openworm.org/].

doi:10.1186/1471-2202-14-S1-P209

Cite this article as: Vella et al.: Integration of predictive-corrective incompressible SPH and Hodgkin-Huxley based models in the OpenWorm in silico model of *C. elegans*. *BMC Neuroscience* 2013 **14** (Suppl 1):P209.

* Correspondence: mv333@cam.ac.uk

¹Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, CB2 3DY, UK

Full list of author information is available at the end of the article