

POSTER PRESENTATION

Open Access

Neural oscillations arising from a linear current with negative conductance

Farzan Nadim^{1,2*}, Yinzheng Guan¹, Jorge Golowasch^{1,2}, Amitabha Bose²

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Introduction

Slow neural oscillations are known to depend on regenerative inward currents. The voltage range in which these currents dominate the neuron's activity is typically limited to the negative-slope conductance range of their IV curve. At low voltages, these currents are not active and at high voltages, outward currents and spike-activity dominate. We find that the contribution of such currents to neural oscillations can be approximated by a linear current with negative slope conductance (I_{NL}). Here we explore the minimal set of requirements to produce neural oscillations in the presence of I_{NL} with the goal of elucidating how other current types shape the properties of oscillations.

Methods

The pacemaker PD neurons in the crab pyloric network are quiescent in the absence of modulatory inputs. We have previously shown that adding I_{NL} with dynamic clamp restores oscillatory activity in these neurons. We now examine whether dynamic clamp I_{NL} can recover oscillations in the absence of spiking (in TTX). We explore the generation of oscillations with I_{NL} using phase-space and bifurcation analysis of simple mathematical models.

Results

We find that adding I_{NL} in the presence of TTX is sufficient to produce oscillatory activity in PD neurons (Fig. A). To understand how oscillations can be generated by adding this simple linear current, we analyzed a 2-variable conductance-based model which includes I_{NL} and an outward current I_K with a single activation variable.

* Correspondence: farzan@njit.edu

¹Department of Biological Sciences, NJIT-Rutgers University, Newark, NJ 07102, USA

Full list of author information is available at the end of the article




Figure 1 A. Addition of I_{NL} to a biological PD neuron in TTX results in oscillatory activity, an effect mimicked by a 2d model containing I_{NL} and I_K . (Ctrl: -50 mV) **B.** The 2d model bifurcation diagram with g_{NL} as parameter shows a limited range of oscillatory activity bound by two Hopf bifurcation points.

We find that this model is able to reproduce the PD oscillations but these oscillations are limited to a small range of g_{NL} values (Fig. B).

The dynamics of a quiescent neuron are locally represented by a stable equilibrium point. Adding I_{NL} destabilizes this equilibrium point. In the absence of a limiting factor, the voltage would grow without bounds. At depolarized voltages, this limiting factor is provided by I_K . Surprisingly, the presence of I_K together with I_{NL} also limits the voltage from escaping in the negative direction and thus results in oscillatory activity (Figure 1B). These oscillations require a delicate balance between I_K and I_{NL} , leading to restrictions in parameter space where rhythmic activity exists. We show that oscillations are readily stabilized in the model by adding a hyperpolarization-activated current such as I_h .

Conclusions

The existence of I_{NL} is sufficient to destabilize a stable equilibrium point around which oscillations can be produced. However, I_{NL} must interact with other membrane currents to be able to bind the voltage both at hyperpolarized and depolarized levels. The existence of these bounds depends on the intrinsic properties of individual neurons which may determine whether or not a neuron can oscillate using I_{NL} . Thus, these results suggest that adding I_{NL} to other pyloric neurons may not lead to oscillations and whether it does will depend on what other ionic currents they express. Our current work involves identifying which pyloric neurons respond to I_{NL} with oscillatory activity and which not, and how that depends on the intrinsic properties of these cells.

Acknowledgements

Supported by NSF1122291.

Author details

¹Department of Biological Sciences, NJIT-Rutgers University, Newark, NJ 07102, USA. ²Department of Mathematical Sciences, NJIT, Newark, NJ 07102, USA.

Published: 8 July 2013

Reference

1. Zhao S, Golowasch J, Nadim F: Pacemaker neuron and network oscillations depend on a neuromodulator-regulated linear current. *Front Behav Neurosci* 2010, **4**:21.

doi:10.1186/1471-2202-14-S1-P32

Cite this article as: Nadim et al.: Neural oscillations arising from a linear current with negative conductance. *BMC Neuroscience* 2013 **14**(Suppl 1):P32.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

