

POSTER PRESENTATION

Open Access


Global network community and non-uniform cell density in the macaque brain

Masanori Shimono

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

The important question, how the global network architecture connecting cortical regions keeps balances between integration and segregation of information processes, have been asked to understand the design of the brain [1,2]. This study aimed to clarify how topological characteristics of such global network architecture relate with physiological characteristics inside of segmented cortical regions in the monkey brain [3]. Especially, I focused on cell densities (densities of neurons or non-neurons) as the representative characteristics of segmented cortical regions [4], and compared the cell densities with network topologies of cortico-cortical fiber tracts [Figure1-A]. To reduce biases in comparisons, I surveyed many topological measures as wide as possible. Total number of evaluated network measures was 27.

As the result, surprisingly, only participation coefficient (PCs) showed significant correlations with cell densities [3]. Although a previous study reported that cell densities significantly change on the anterior-posterior coordinate [5], spatial coordinates did not correlate significantly with participation coefficients. Participation coefficient is the topological measure evaluating how often each node (segmented brain region) connects to other nodes locating different communities (modules). The modules, which detected based on a computational criterion [3], corresponded with visual, somatosensory, auditory, and two associative modules [Figure1-C]. The associative modules simultaneously showed low neuron density and high participation coefficient, which means there are diverse connections with different modules. These findings led us to



Correspondence: mshimono@indiana.edu
Department of Physics, University of Indiana, Bloomington, IN, 47405, USA

the conclusion that the brain is designed for achieving integrative information process at associative brain regions by sacrificing number of elements (neurons).

Acknowledgements

M.S. is grateful to Yasuo Kawaguchi, John M. Beggs, and Olaf Sporns for their important discussions. This study was supported by a Grant-in-Aid for JSPS Fellows for Research Abroad.

Published: 21 July 2014

References

1. Chen Y, Wang S, Hilgetag CC, Zhou C: Trade-off between Multiple Constraints Enables Simultaneous Formation of Modules and Hubs in Neural. *PLoS computational biology* 2013, **9**(3):e1002937.
2. Sporns O: Network attributes for segregation and integration in the human brain. *Current opinion in neurobiology* 2013.
3. Shimono M: Non-uniformity of cell density and networks in the monkey brain. *Scientific reports* 2013, **3**.
4. Collins CE, Airey DC, Young NA, Leitch DB, Kaas JH: Neuron densities vary across and within cortical areas in primates. *Proc Natl Acad Sci USA* 2010, **107**(36):15927-15932.
5. Cahalane DJ, Charvet CJ, Finlay BL: Systematic, balancing gradients in neuron density and number across the primate isocortex. *Frontiers in Neuroanatomy* 2012, **6**:28 .

doi:10.1186/1471-2202-15-S1-P100

Cite this article as: Shimono: Global network community and non-uniform cell density in the macaque brain. *BMC Neuroscience* 2014 **15** (Suppl 1):P100.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

