

POSTER PRESENTATION

Open Access


Bursting suppression in propofol-induced general anesthesia as bi-stability in a non-linear neural mass model

Pedro Garcia-Rodriguez*, Axel Hutt

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Bursting activity suppression, a phenomenon characterized by sequences of alternating quiescent ('down') and bursting ('up') states, is ubiquitously present during

deep sedation in general anesthesia and can be observed in LFPs and in EEG recordings as well [1,2]. However, the dynamical principles underlying such phenomena



* Correspondence: pedro.garcia-rodriguez@inria.fr
INRIA CR Nancy-Grand Est, Equipe NeuroSys, France

are unclear. Prevailing theoretical approaches to this problem suggest that such bi-stable dynamics could be explained by the existence of notable non-linearities in the neuronal membrane potential as a function of the anesthetic agent, with frequent noise-induced transitions between two stable (attracting) branches. Nevertheless, the mathematical tractability of those models is rather limited due to the dimensionality (2D) of the phase-space (see Figure 1). It has been shown for a neural-mass version of the model proposed in [3], that the 2D-trajectory likely escapes to the other attractor through a limited section of the saddle-line separating the attractions basins [4]. This prediction is here supported by numerical simulations, permitting the simplification of the dynamics to a simpler two-state trace. Relevant statistics can then be extracted, including histograms of the duration of the visit in each state that may support future analytical calculations of the EEG power spectrum following renewal theory [5]. Interestingly, it is found that only the average waiting time in the 'up' states is a non-linear decreasing function of p , differently to the residence time in the 'down' state that behaves linearly in p and in the synaptic time-scale ($1/\alpha$) direction as well.

Published: 21 July 2014

References

1. Ferron JF, et al: *J. Neuroscience* 2009, **29**(31):9850-9860.
2. Wilson M, et al: *J Biological Physics* 2010, **36**:245-259.
3. Hutt A, Longtin A: *Cognitive Neurodynamics* 2009, **4**(1):37-59.
4. Rodriguez P, Hutt A: *Bernstein Conference* 2013.
5. Stratonovich RL: *Topics in the theory of random noise:Volume I*, Ch. 6. *Gordon and Breach* 1962.

doi:10.1186/1471-2202-15-S1-P139

Cite this article as: Garcia-Rodriguez and Hutt: Bursting suppression in propofol-induced general anesthesia as bi-stability in a non-linear neural mass model. *BMC Neuroscience* 2014 **15**(Suppl 1):P139.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

