

POSTER PRESENTATION

Open Access

Afferent-hair cell connectivity as a possible source of spike train irregularity in turtle vestibular bouton afferents

William R Holmes*, Janice A Huwe, Michael H Rowe, Ellengene H Peterson

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Vestibular afferents have been categorized as “regular” or “irregular” depending on the variability of their spike trains [1,2]. It is thought that the difference in regularity among afferents results from differences in the types of voltage-dependent conductances present in the cell. For example afferents may differ in their complements of potassium A, D, and M currents or in currents that produce a small vs. prominent afterhyperpolarization (AHP) [3-5]. Because bouton afferents from turtle utricle vary widely in morphology depending on their location in the utricle, we sought to determine if afferent morphology might also play a role in spike train irregularity. Although we found that morphology itself is not a factor in determining spike train variability, we did find that spike train variability can be affected by afferent/hair cell connectivity.

Morphological reconstructions of bouton afferents from the lateral extrastricular (LES), striola, juxtastricola and medial extrastricular (MES) regions of turtle utricle were done with NeuroLucida. Models based on these reconstructions were constructed for 16 afferents. Voltage-dependent conductance distributions in different regions (axon, soma, terminal arbor, varicosities, nodes, myelin) were identical in each cell model. Given the locations of synapses in the terminal arbor and the density of hair cells in each region, we were able to estimate the number of synapses formed between individual hair cells and an afferent. In the LES, striola and juxtastricola most synapses on an afferent were formed with unique hair cells, whereas in the MES single hair cells more often made multiple synapses with an afferent. To determine the effect of this difference in connectivity we modeled Poisson release from hair cells to synapses in the terminal

arbor to generate spike trains. The coefficient of variation (cv) as a function of interspike interval (ISI) was calculated to estimate spike train variability with different average release rates.

We found that cv as a function of ISI was nearly identical for LES, striola and juxtastricola afferents, but was the same for MES afferents only if connectivity in the MES was assumed to be one afferent synapse per hair cell. However, cv was much larger when the MES afferents were modeled with multiple synapses between an afferent and a hair cell, as calculated from synaptic location and hair cell density. To see if we could make an LES afferent more irregular, we modified connectivity to have more multiple synapses between individual hair cells and the afferent. As predicted, cv increased.

These results suggest that afferent-hair cell connectivity may be a source of spike train irregularity. Given the large number of multiple contacts between a calyx and its presynaptic hair cell, we expect that connectivity contributes to spike train variability in calyx-bearing (calyx and dimorph) afferents.

Published: 21 July 2014

References

1. Baird RA, Desmadryl G, Fernández C, Goldberg JM: The vestibular nerve of the chinchilla. II. Relation between afferent response properties and peripheral innervation patterns in the semicircular canals. *J Neurophysiol* 1988, **60**:182-203.
2. Goldberg JM, Desmadryl G, Baird RA, Fernández C: The vestibular nerve of the chinchilla. IV. Discharge properties of utricular afferents. *J Neurophysiol* 1990, **63**:781-790.
3. Goldberg JM, Smith CE, Fernández C: Relation between discharge regularity and responses to externally applied galvanic currents in vestibular nerve afferents of the squirrel monkey. *J Neurophysiol* 1984, **51**:1236-1256.

* Correspondence: holmes@ohio.edu

Department of Biological Sciences, Neuroscience Program, Ohio University, Athens, OH 45701, USA

4. Kalluri R, Xue J, Eatock RA: Ion channels set spike timing regularity of mammalian vestibular afferent neurons. *J Neurophysiol* 2010, **104**:2034-2051.
5. Eatock RA, Xue J, Kalluri R: Ion channels in mammalian vestibular afferents may set regularity of firing. *J Exp Biol* 2008, **211**:1764-1774.

doi:10.1186/1471-2202-15-S1-P69

Cite this article as: Holmes *et al.*: Afferent-hair cell connectivity as a possible source of spike train irregularity in turtle vestibular bouton afferents. *BMC Neuroscience* 2014 **15**(Suppl 1):P69.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

