

POSTER PRESENTATION

Open Access

# Pattern recognition of Hodgkin-Huxley equations by auto-regressive Laguerre Volterra network

Kunling Geng<sup>1,2\*</sup>, Vasilis Z Marmarelis<sup>1,2</sup>

From 24th Annual Computational Neuroscience Meeting: CNS\*2015  
Prague, Czech Republic. 18-23 July 2015

A nonparametric, data-driven nonlinear auto-regressive Volterra (NARV) [1] model has been successfully applied for capturing the dynamics in the generation of action potentials, which is classically modeled by Hodgkin-Huxley (H-H) equations. However, the compactness still need to be improved for further interpretations. Therefore, we propose a novel Auto-regressive Sparse Laguerre Volterra Network (ASLVN) model (shown in Figure 1A), which is developed from traditional Laguerre Volterra Network (LVN) and principal dynamic mode (PDM) framework [2].

We adopt stochastic global optimization algorithm Simulated Annealing [3] to train the ASLVN instead of Back-propagation method [2] to avoid local minima and convergence problems. We also use lasso regularization [4] to enhance the sparsity of the network and prune redundant branches for parsimony. The prediction results are shown in Fig.1B, it can be seen that the exogenous output  $z^{(1)}$  represents the subthreshold dynamics in phase III, and the autoregressive output  $z^{(2)}$  dominates in the spike shape in phase I, and the cross term output  $z^{(x)}$  helps to maintain the refractory period by cancelling the effect of  $z^{(1)}$  in phase II and we also observe that refractory inhibition effect decays after initiation of AP, which explains the absolute refractory period and relative refractory period in physiology.

#### Authors' details

<sup>1</sup>Biomedical Engineering Department, University of Southern California, Los Angeles, CA 90089, USA. <sup>2</sup>Biomedical Simulations Resource, Los Angeles, CA 90089, USA.

Published: 18 December 2015

\* Correspondence: kgeng@usc.edu

<sup>1</sup>Biomedical Engineering Department, University of Southern California, Los Angeles, CA 90089, USA

Full list of author information is available at the end of the article

#### References

1. Eikenberry SE, Marmarelis VZ: A nonlinear autoregressive Volterra model of the Hodgkin-Huxley equations. *Journal of computational neuroscience* 2013, **34**(1):163-183.
2. Marmarelis VZ: Nonlinear dynamic modeling of physiological systems. *John Wiley & Sons* 2004, **10**.
3. Kirkpatrick S: Optimization by simulated annealing. *science* 1983, **220**(4598):671-680.
4. Tibshirani R: Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B (Methodological)* 1996, **267**-288.

doi:10.1186/1471-2202-16-S1-P156

Cite this article as: Geng and Marmarelis: Pattern recognition of Hodgkin-Huxley equations by auto-regressive Laguerre Volterra network. *BMC Neuroscience* 2015 **16**(Suppl 1):P156.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



