

Poster presentation

Open Access

Measuring spike train synchrony and reliability

Thomas Kreuz^{*1}, Julie S Haas², Alice Morelli³, Henry DI Abarbanel^{2,4} and Antonio Politi¹

Address: ¹Istituto dei Sistemi Complessi – CNR, Sesto Fiorentino, Italy, ²Institute for Nonlinear Sciences, University of California, San Diego, CA, USA, ³Istituto Nazionale di Ottica Applicata, Firenze, Italy and ⁴Department of Physics and Marine Physical Laboratory (Scripps Institution of Oceanography), University of California, San Diego, CA, USA

Email: Thomas Kreuz* - thomas.kreuz@fi.isc.cnr.it

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, **8**(Suppl 2):P79 doi:10.1186/1471-2202-8-S2-P79

© 2007 Kreuz et al; licensee BioMed Central Ltd.

Estimating the degree of synchrony or reliability between two or more spike trains is a frequent task in both experimental and computational neuroscience. In recent years, many different methods have been proposed that typically compare the timing of spikes on a certain time scale to be fixed beforehand. In this study [1], we propose the ISI-distance, a simple complementary approach that extracts information from the interspike intervals by evaluating the ratio of the instantaneous frequencies. The method is parameter free, time scale independent and easy to visualize as illustrated by an application to real neuronal spike trains obtained *in vitro* from rat slices (cf. [2]). We compare the method with six existing approaches (two spike train metrics [3,4], a correlation measure [2,5], a similarity measure [6], and event synchronization [7]) using spike trains extracted from a simulated Hindmarsh-Rose network [8]. In this comparison the ISI-distance performs as well as the best time-scale-optimized measure based on spike timing, without requiring an externally determined time scale for interaction or comparison.

Acknowledgements

TK has been supported by the Marie Curie Individual Intra-European Fellowship "DEAN", project No 011434. JSH acknowledges financial support by the San Diego Foundation.

References

1. Kreuz T, Haas JS, Morelli A, Abarbanel HDI, Politi A: **Measuring spike train synchronization.** [<http://arxiv.org/abs/physics/0701261>].
2. Haas JS, White JA: **Frequency selectivity of layer II stellate cells in the medial entorhinal cortex.** *J Neurophysiol* 2002, **88**:2422-2429.
3. Victor J, Purpura K: **Nature and precision of temporal coding in visual cortex: A metric-space analysis.** *J Neurophysiol* 1996, **76**:1310.
4. van Rossum MCW: **A novel spike distance.** *Neural Computation* 2001, **13**:751.
5. Schreiber S, Fellous JM, Whitmer JH, Tiesinga PHE, Sejnowski TJ: **A new correlation-based measure of spike timing reliability.** *Neurocomputing* 2003, **52**:925.
6. Hunter JD, Milton G: **Amplitude and frequency dependence of spike timing: implications for dynamic regulation.** *J Neurophysiol* 2003, **90**:387.
7. Quiroga R, Kreuz T, Grassberger P: **Event synchronization: A simple and fast method to measure synchronicity and time delay patterns.** *Phys Rev E* 2002, **66**:041904.
8. Morelli A, Grotto RL, Arechi FT: **Neural coding for the retrieval of multiple memory patterns.** *Biosystems* 2006, **86**:100.