Review

Open Access

Behavioral genetics and taste

John D Boughter Jr*¹ and Alexander A Bachmanov²

Address: ¹Department of Anatomy & Neurobiology, University of Tennessee Health Science Center, Memphis, TN 38163, USA and ²Monell Chemical Senses Center, 3500 Market Street, Philadelphia, PA 19104-3308, USA

Email: John D Boughter* - jboughter@utmem.edu; Alexander A Bachmanov - bachmanov@monell.org

* Corresponding author

Published: 18 September 2007

BMC Neuroscience 2007, 8(Suppl 3):S3 doi:10.1186/1471-2202-8-S3-S3

This article is available from: <http://www.biomedcentral.com/1471-2202/8/S3/S3>

© 2007 Boughter and Bachmanov; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This review focuses on behavioral genetic studies of sweet, umami, bitter and salt taste responses in mammals. Studies involving mouse inbred strain comparisons and genetic analyses, and their impact on elucidation of taste receptors and transduction mechanisms are discussed. Finally, the effect of genetic variation in taste responsiveness on complex traits such as drug intake is considered. Recent advances in development of genomic resources make behavioral genetics a powerful approach for understanding mechanisms of taste.

Introduction

The chemical senses, gustation and olfaction, have provided a candidate media for investigation of the effects of major genes on behavior, owing to considerable natural variation for both senses among individuals or strains of mice and rats. Genetic analysis of behavioural responses to bitter and sweet taste stimuli in mice facilitated the recent discovery of receptor gene families for bitter, sweet and amino acid taste (for recent reviews, see [1-4]). The function and specificity of taste receptors continues to be elucidated based on not only *in vitro* expression assays and transgenic approaches, but also based on studies of gene variation and evolution (e.g. [5-8]).

Taste, beyond its purely sensory function, is also inextricably linked to a larger set of behaviors. The gustatory system is anatomically located at the beginning of the alimentary canal, and as such is a key player in homeostatic systems dealing with nutrient and fluid intake. Taste and feeding are linked in terms of reciprocal connectivity between brain centers (e.g. [9,10]). The consumption of sweeteners, for example, often involves a complex integration of the peripheral sensory system, the central nervous

system and post-ingestive events. Sweet taste detection in mammals has been shown to be heavily modulated by post-ingestive feedback, including neural or hormonal factors (e.g. [11-13]). This feedback extends to the level of the taste receptor cells and afferent nerves [14,15]. Nutritive sweeteners provide post-ingestive stimulation [16], probably due to their calories. These interactive mechanisms of consummatory responses to sweeteners imply that ingestion is a complex behavior, and is likely to be determined by multiple genes.

Although studies of non-mammalian organisms such as fruit flies hold tremendous potential for elucidating genetic basis of taste behavior and physiology [17,18] this review will focus on studies of taste genetics in mammals, primarily in mice, and to a lesser extent in rats and humans. In particular, we will concentrate on the characterization of phenotypic diversity among inbred strains of mice, as this naturally-occurring variation has proved crucial for genetic analysis and the eventual identification of G-protein coupled taste receptors. Hence, the review is arranged by taste quality, including those characterized by humans as tasting sweet, umami (savory), bitter and salty.

Although candidate sour taste receptor molecules have been recently identified [19,20], less progress has been made regarding genetic approaches to sour (acid) taste, and this quality is not discussed here.

Measuring taste behavior

Taste perception involves several aspects, such as intensity, quality and hedonic value of the taste sensation. In humans, these aspects of taste perception are usually assessed using verbal information, e.g., by plotting sensation intensity on a scale with verbal descriptors or by reporting a difference between samples. Early studies of human genetic variation of taste sensitivity concentrated on either responses to a single concentration of stimulus (i.e., tasters vs. non-tasters of phenylthiocarbamide (PTC)) or measurement of absolute thresholds [21]. A more expansive view of taste ability, and therefore phenotypic differences, may be gained from analyses of the perceived taste intensity [22,23].

Assessment of taste perception in non-human animals relies on recording of behavior elicited by taste stimuli using a number of different techniques [24]. Each of these techniques characterizes some, but not all aspects of taste perception. For example, hedonics of taste sensation can be assessed based on consummatory responses in naive animals, but assessing taste quality involves conditioning of animals. Many of these techniques have been adapted for use in mice and can be used in genetic experiments (Table 1).

Traditionally, behavior genetic studies of taste in the mouse have been conducted using two-bottle intake tests, e.g. [25-28]. These tests simply measure the consumption of a taste stimulus over a 24- or 48-h period, relative to the consumption of water. Sweet-tasting stimuli are generally preferred to water, whereas bitter-tasting stimuli are avoided. Such tests, while amenable for testing large cohorts of mice required for genetic analysis, do not easily allow for the dissociation of sensory ability from post-ingestive factors such as satiety or toxicity. Brief-access tests minimize post-ingestive cues by restricting trial lengths to short durations (< 30 s). The dependent measure is number of licks taken from a particular stimulus, typically expressed as a ratio (licks to stimulus/licks to water). Such tests have been effectively used for high-throughput screening of taste function in mouse populations, especially for compounds with an aversive taste [29,30].

Other tests of taste function in the mouse include measurements of threshold detection or the ability to discriminate between two stimuli [31,32]. Conditioned taste aversion (CTA) is a commonly used paradigm to estimate perceptual similarity or dissimilarity between stimuli. The

presentation of a novel taste stimulus is paired with a stimulus that produces temporary gastric distress, usually an intraperitoneal dose of LiCl. After pairing, the animal will avoid consumption of the conditioned taste stimulus, and this CTA will generalize to stimuli with similar perceptual qualities.

Finally, electrophysiological recordings of activity in the afferent gustatory nerves often corroborate behavioral taste tests in non-human animals. The major gustatory nerves are the chorda tympani (CT) branch of the facial (VII) nerve and the glossopharyngeal (IX) nerve. The CT predominantly innervates taste buds on the anterior tongue, whereas the IXth nerve innervates taste buds contained in trenches on the posterior tongue. Studies of gustatory nerve activity help to elucidate whether genetic effects on taste perception have peripheral or central origin.

Genetic analysis of taste behavior

In genetic terms, the taste response of an individual represents a phenotype, i.e., an observable property of an organism produced by genetic and environmental factors. A typical goal of genetic studies is to characterize the genetic factors contributing to the phenotype. This is usually achieved by analyzing populations of related individuals, such as twins or families in humans, or crosses between inbred strains of mice or rats. An initial goal of the genetic analysis is to establish whether the phenotypic variation among individuals has a genetic component (i.e., is heritable). Once heritability is confirmed, chromosomal localization of genes responsible for the heritable component of the phenotype can be found using linkage analysis. The linkage analysis involves assessment of association between genetic markers distributed throughout the genome and the phenotype. Some linkage studies in rodents use recombinant inbred strains, which are produced by intercrossing two parental inbred strains, obtaining the second generation of hybrids (F_2), and subsequent inbreeding (brother $\times$ sister mating) of multiple strains. After a genetic locus affecting a phenotype is mapped to a particular chromosome (Chr), subsequent studies can identify the molecular nature of the corresponding gene. Because gene identification is based primarily on the chromosomal position of the gene, this approach is called positional cloning.

Sweet and umami taste

Genetic variation in sweet taste responsiveness

Taste sensitivity to sweeteners varies among individual humans, and this variation may be determined genetically [33-42]. Prominent genetic differences in taste responses to sweeteners also exist among inbred strains of mice. These differences were shown using different experimental techniques and a variety of sweeteners (such as sucrose,

Table 1: Methods to characterize taste in non-human animals

Information	Method			
	Long-term two-bottle preference	Brief-access licking responses	CTA	Electrophysiological recording of taste-evoked activity in gustatory nerves
Sensitivity (threshold)	+/- ^a	+/- ^a	+	+
Intensity	+	+	-	+
Hedonic (pleasant or unpleasant) properties	+	+	-	-
Taste quality	-	-	+	+/- ^b
Specificity	Can be influenced by post-ingestive effects	Minimizes postingestive effects; can be influenced by water restriction	Can be influenced by non-taste sensory properties (e.g., odor), learning ability or sensitivity to an unconditioned stimulus	Directly assesses peripheral neural response

+, test is primarily designed to assess a particular aspect of taste perception (e.g., intensity or quality); +/-, limited information regarding the aspect of taste perception can be derived from test results; -, test is not informative regarding the aspect of taste perception. ^a Response thresholds of naive (not conditioned) animals are often higher than taste recognition thresholds in conditioned animals. ^b Taste quality can be assessed using analysis of single-fiber activity, but not using integrated whole-nerve recordings. All types of tests have been used in genetic experiments, but two-bottle preference tests have the highest throughput of phenotyping.

glucose, dulcin, saccharin, acesulfame, glycine, D-phenylalanine and L-glutamine). Mice from different strains vary in taste responses to sweeteners assessed using long-term preference tests [26,27,43-53], single-bottle tests [54], brief-access tests based on lick recording [55,56], taste detection thresholds [31], CTA generalization [57], and responses of gustatory nerves [58-60]. These studies have shown that responses to many of these sweeteners (e.g., sucrose, glucose, dulcin, saccharin and acesulfame) closely correlate among mouse strains, suggesting a common genetic basis for sweet taste. However, responses to some sweet-tasting amino acids display somewhat different patterns of strain differences. Strain differences in consummatory responses to sweeteners have also been reported for rats [61-63] and hamsters [64].

A few mouse strains with large differences in sweet taste responses were used to produce crosses for analyses of genetic and physiological mechanisms underlying behavioral responses to sweeteners. The most detailed physiological analysis was conducted using mice from the C57BL/6 (B6) strain with high sweetener preferences and mice from the 129 strains with low sweetener preferences. Compared with 129 mice, B6 mice had higher preferences for a large number of sweeteners, including sugars (sucrose and maltose), sweet-tasting amino acids (glycine, D-phenylalanine, D-tryptophan, L-proline and L-glutamine), and several but not all non-caloric sweeteners (saccharin, acesulfame, dulcin, sucralose and SC-45647) [46,50-52,65]. This phenotypic difference is specific to sweet-taste processing, and is not due to a generalized difference in taste responsiveness or differences in appetite [66,67].

Differences between B6 and 129 mice in preference for a sweet-tasting amino acid glycine [65] appear to depend on mechanisms distinct from those affecting responses to many other sweeteners. Both B6 and 129 mice generalized a CTA between glycine and several other sweeteners, demonstrating that they perceive the sucrose-like taste of glycine. Thus, the lack of a strong glycine preference by 129 mice cannot be explained by their inability to perceive its sweetness [68]. Despite differences in glycine intakes and preferences, CT responses to glycine are similar in mice from both strains [58]. Neither behavioral nor neural responses to glycine are influenced by the *Tas1r3* genotype [31,69], suggesting that variation in taste responses to glycine depends on other genes.

Some genetic analyses of sweetener consumption by mice yielded evidence that it is influenced by a single locus, named *Sac* (saccharin preference) [46,50,70,71], whereas other experiments indicated that more than one gene is involved [47,50,51,72,73]. The apparent discrepancy on whether single-gene or multi-gene model better describes genetic variation in sweetener preferences is likely due to use of different progenitor strains and types of mapping panels, different sweetener solutions tested, and different quantitative analyses used in these studies.

The saccharin preference (*Sac*) locus and the *Tas1r3* gene
Using long-term two-bottle tests, Fuller [70] demonstrated that differences in saccharin preferences between the B6 and DBA/2J inbred strains largely depend on a single locus, *Sac*, with a dominant *Sac^b* allele present in the B6 strain associated with higher saccharin preference, and a recessive *Sac^d* allele present in the DBA/2J strain associ-

ated with lower saccharin preference. Subsequent studies confirmed this finding in the BXD recombinant inbred strains, and in crosses between the B6 and DBA/2 or between the B6 and 129 strains [46,50,71,72,74,75]. In addition to sweetener preferences, the *Sac* genotype influences the afferent responses of gustatory nerves to sweeteners [75,76], which indicated that the *Sac* gene is involved in peripheral taste transduction and may encode a sweet taste receptor.

The *Sac* locus has been mapped to the subtelomeric region of mouse Chr 4 [50,72,74-76] (see also Figure 1). A positional cloning study at the Monell Chemical Senses Center showed that the *Sac* locus corresponds to the *Tas1r3* gene [77,78]. This study involved a high-resolution linkage analysis of a B6 × 129 F₂ intercross, the marker-assisted selection of a 129.B6-*Sac* congenic strain, physical mapping that involved construction of a contig of bacterial artificial chromosome (BAC) clones, BAC sequencing, and sequence analysis of candidate genes. One of the genes within the critical interval of the *Sac* locus was a G protein-coupled receptor gene, *Tas1r3* (taste receptor, type 1, member 3). Based on the effects of the *Sac* genotype on peripheral sweet taste responsiveness [75,76], and on involvement of a G protein-coupled mechanism in sweet taste transduction [79], *Tas1r3* was selected as the most likely candidate for the *Sac* locus (Figure 1).

The *Tas1r3* gene contains 6 coding exons and is translated into an 858-amino acid protein, T1R3, with a predicted secondary structure that includes seven transmembrane domains and a large hydrophilic extracellular N-terminus. This structure is typical of the class C G protein-coupled receptor family, which includes the metabotropic glutamate and extracellular calcium-sensing receptors. T1R3 belongs to a small family of G protein coupled receptors, T1R, which also includes the T1R1 and T1R2 proteins. The three mouse T1R genes are located on distal Chr 4 in the order: *Tas1r2* (70.0 cM, or 139 Mb, NCBI Build 36) – *Tas1r1* (81.5 cM, or 151 Mb) – *Tas1r3* (83.0 cM, or 155 Mb). The *Tas1r1* and *Tas1r2* genes were excluded as candidates for the *Sac* locus based on their more proximal chromosomal location [76,80,81]. All three T1R members are expressed in taste receptor cells [53,80-85]. When T1R3 is co-expressed in a heterologous system with T1R2, it functions as a sweet receptor [85-87], but T1R3 may also function as a low-affinity sugar receptor alone, probably as a homodimer [88]. *Tas1r3* has a human ortholog, *TAS1R3*, residing in a region of conserved synteny in the short arm of human Chr 1 (1p36) [89].

If *Tas1r3* is identical to *Sac*, substitution of *Tas1r3* alleles must result in phenotypical changes attributed to the *Sac* locus. Introgression of the 194-kb chromosomal fragment containing the *Tas1r3* allele from the high-sweetener pre-

ferring B6 strain onto the genetic background of the 129 strain fully rescued its low sweetener preference phenotype: sweetener intake of the congenic mice was as high as that of mice from the donor B6 strain [77] (Figure 1). Equivalent phenotype rescue results were obtained in a transgenic experiment [85]. These data demonstrate that substitution of *Tas1r3* alleles results in behavioral changes attributed to the *Sac* locus and therefore provides a proof that *Tas1r3* is identical to *Sac*, and that the T1R3 receptor responds to sweeteners. Further evidence for identity of the *Sac* locus and *Tas1r3* gene was obtained in studies of mice with targeted mutations of the *Tas1r3* gene, which were found to be deficient in taste responses to sweeteners [88,90].

Identity of *Sac* and *Tas1r3* implies that there must be *Tas1r3* polymorphisms, resulting in variation of sweet taste responses attributed to allelic variants of the *Sac* locus. Although several candidate functional polymorphisms were proposed in studies that identified the *Tas1r3* gene [80,81,83-85], these studies lacked a proper quantitative analyses of gene-phenotype associations.

Reed et al [53] conducted a comprehensive quantitative analysis of the *Tas1r3* sequence variants associated with saccharin preference using 30 genealogically diverse inbred mouse strains. Genomic sequences including *Tas1r3* exons, introns, upstream and downstream regions were examined, so that polymorphisms affecting amino acid composition or potential regulatory regions could be detected. The strongest association with saccharin preference was found for a haplotype including three sites: nucleotide (nt) -791 (3 bp insertion/deletion), nt +135 (Ser45Ser), and nt +179 (Ile60Thr) (Figure 2). Lack of differences in the *Tas1r3* gene expression in the taste tissues of mice with different *Tas1r3* haplotypes suggested that the polymorphisms that do not change amino acid sequence of the T1R3 protein (nt -791 and nt +135) are unlikely to affect receptor function. Therefore, the amino acid substitution of isoleucine to threonine at position 60 (Ile60Thr) was predicted to be a functional polymorphism [53]. This prediction was subsequently confirmed in an *in vitro* study showing that a corresponding site-directed mutation changes binding affinity of the T1R3 protein to several sweeteners [91].

Ligand specificity of the T1R3 receptor

In vitro studies of T1R proteins have indicated that they function as broad-spectrum sweet and umami receptors (Table 2) [85,87,92]. Several aspects of these *in vitro* studies emphasize importance of an *in vivo* approach to characterize ligand-receptor interactions. First, discrepancies between results obtained using different expression systems leave open a question of whether responsiveness or unresponsiveness to a particular sweetener reflects *in vivo*

Figure 1

Positional identification of the Sac (saccharin preference) locus. **a.** Linkage map of mouse distal Chr 4 based on data from the B6 × 129 F₂ intercross. The X axis shows distances between markers in recombination units (cM). The Y axis shows the logarithm of the odds ratio (LOD) scores for sucrose and saccharin consumption. The LOD score peaks (indicated by black triangles) and confidence intervals (solid horizontal line for sucrose, 4.5 cM, and dotted horizontal line for saccharin, 5.3 cM) define the genomic region of the Sac locus. **b.** Average daily 17 mM saccharin consumption by mice from parental 129 and B6 strains (left), B6 × 129 F₂ hybrids (center), and congenic 129.B6-Sac mice (right) in 96-hr two-bottle tests with water (means ± SE). *Tas1r3* genotypes of the F₂ and congenic mice and mouse numbers are indicated on the bars. Differences between parental strains and among the F₂ and congenic genotypes were significant ($p < 0.0001$, ANOVA). F₂ and congenic B6 homozygotes and heterozygotes for *Tas1r3* did not differ from each other, and had higher saccharin intakes compared with 129 homozygotes ($p < 0.0001$, post hoc tests). **c.** Linkage map of the Sac-containing region defined based on the size of the donor fragment in the 129.B6-Sac congenic strain (black box). Distances between markers were estimated based on the B6 × 129 F₂ intercross (see panel a). **d.** A contig of bacterial artificial chromosome (BAC) clones and physical map of the Sac region. BAC clones are represented by horizontal lines. Dots indicate marker content of the BAC clones. **e.** Genes within the Sac-containing interval. Filled areas indicate predicted genes. Arrows indicate the predicted direction of transcription. Figure reproduced with permission from [253].

Figure 2
Preference for 1.6 mM saccharin by mice from inbred strains with different *Tas1r3* genotypes at the T/C variant site at nucleotide position +179 (relative to the first nucleotide in the ATG start codon of the *Tas1r3* gene). This polymorphism results in amino acid substitution of isoleucine to threonine at position 60 (I60T), in the extracellular N-terminus of the predicted T1R3 protein. Closed circles denote means for C57BL/6, C57L/J, CAST/Ei, CE/J, FVB/NJ, I/LnJ, IS/Cam, KK/HIJ, NOD/LtJ, NZB/BINJ, P/J, RBF/DnJ, SEA/GnJ, SJL/J, SM/J, SPRET/Ei, ST/bj and SWR/J strains with +179 T genotype. Mice from these strains strongly preferred saccharin (average preference score $88 \pm 2\%$, Mean $\pm$ SE; $n = 18$). Open circles show means for 129P3/J, A/J, AKR/J, BALB/cByJ, BUB/BnJ, C3H/HeJ, CBA/J, DBA/2J, LP/J, PL/J, RF/J and RIIIS/J strains with +179 C genotype. Mice from these strains were indifferent to or only weakly preferred saccharin (average preference score $59 \pm 3\%$, $n = 12$; $p = 0.0000000012$, t-test). Despite the strong phenotypical effect of the *Tas1r3* genotype, there is also substantial variation in saccharin preference within each genotype group. As a result, *Tas1r3* genotype explains only 78% of genetic variation in saccharin preferences among the inbred strains; the remaining 22% of genetic variance is attributed to the effect of other genes. Adapted with permission from [53].

sensitivity of the receptor, or is an artifact of the *in vitro* system. Second, responses of the heterologously expressed mouse receptors to amino acids are not always consistent with mouse behavioral responses to these stimuli. For

example, sweet L-proline and L-threonine did not activate the T1R2 and T1R3 combination, but instead activated the T1R1 and T1R3 combination, which also responded to some umami-tasting and bitter (e.g., L-phenylalanine) compounds [87].

The *in vivo* approach to characterize ligand specificity of the T1R3 receptor is based on an assumption that if a response to a compound is affected by *Tas1r3* genotype, then this compound activates a receptor involving T1R3. Several earlier genetic mapping studies have shown that in addition to saccharin preferences, the *Sac* locus also affects consumption of sucrose, acesulfame [46,50,74,75], and ethanol, which has a sweet taste component [93], and also chorda tympani responses to saccharin and sucrose [75,76]. Inoue et al. [69] examined behavioral and neural responses to a larger set of sweeteners in the F_2 hybrids between the B6 and 129 strains. They found that the *Tas1r3* genotype affected consumption of the sweeteners sucrose, saccharin and D-phenylalanine, but not glycine. For CT responses, significant linkages to *Tas1r3* were found for the sweeteners sucrose, saccharin, D-phenylalanine, D-tryptophan and SC-45647, but not glycine, L-proline, L-alanine or L-glutamine. No linkages to the *Tas1r3* chromosomal region were detected for behavioral or neural responses to non-sweet quinine, citric acid, HCl, NaCl, KCl, monosodium glutamate (MSG), inosine 5'-monophosphate (IMP) or ammonium glutamate. Thus, allelic variation of the *Tas1r3* gene affects taste responses to many but not all sweeteners, suggesting that a wide variety of sweeteners can activate a receptor involving T1R3. Lack of the effect of the *Tas1r3* genotype on taste responses to some sweeteners, such as glycine, can be explained by several possible mechanisms: (i) sweetener binding to the T1R3 receptor at a site that is not affected by the polymorphic variants; (ii) binding to the T1R2 receptor; (iii) existence of another sweet taste receptor binding these sweeteners. These mechanisms can be examined using mice with targeted mutations of the *Tas1r* genes, but there are currently no data on taste responses to glycine in *Tas1r* knockout mice.

Other genes involved in sweet taste responses

Multigenic inheritance of sweetener preferences was shown in a number of studies [47,50,51,72,73]. Accordingly, several lines of evidence indicated that allelic variation of the mouse *Tas1r3* locus does not account for all the genetically determined differences in sweetener preferences. Analysis of multiple inbred mouse strains has shown that the *Tas1r3* genotype explains only 78% of genetic variation in saccharin preference [53] (Figure 2). In the B6 $\times$ 129 F_2 cross, the *Tas1r3* genotype explained 64 – 96% of genetic variation in preference scores for different sweeteners, 10 – 35% of genetic variation in sweetener intakes, and 37 – 92% of genetic variation in CT responses

Table 2: Deorphanized taste receptors

Ligands	Receptors	
	Human	Mouse
Sugars, sweeteners, D-amino acids ¹	hTIR2 + hTIR3	mTIR2 + mTIR3
Umami stimuli and L-amino acids	hTIR1 + hTIR3	mTIR1 + mTIR3
Beta-glucopyranosides	hT2R16	
PROP	hT2R4 ²	mT2R8 ²
Structurally diverse set of bitter stimuli	hT2R14	
Strychnine	hT2R10	
Saccharin, acesulfame K	hT2R44, hT2R43	
PTC	hT2R38	
Denatonium	hT2R4, hT2R44	mT2R8
Cycloheximide		mT2R5

The ligand specificities have been determined predominantly by *in vitro* studies, but also through the use of knock-out or transgenic mice, as well as other approaches examining allelic variance. The sweet and amino acid receptors have broad ligand specificity. The TIR2 + TIR3 heterodimer responds to a broad array of sweet-tasting stimuli in both human and mouse [85–87]. TIR1 + TIR3 responds to umami stimuli in humans and to a wider range of stimuli including a variety of L-amino acids in mice [86, 87]. A TIR3 homodimer may respond to sugars at higher concentrations [88, 90]. In contrast, most of the bitter receptors have narrow ligand specificities [111, 112, 125, 250, 251]. However, hT2R14 has been shown to respond to a broad group of bitter compounds, including plant neurotoxins [252].

¹Excepting D-glutamate and D-aspartate.

²Only high concentrations of PROP were found to activate this receptor *in vitro*.

to sweeteners [69,75]. Responses to sweeteners in brief-access tests differ among mouse strains but do not seem to be associated with *Tas1r3* alleles [55]. Thus, a substantial part of the genetic variation in taste responses to sweeteners among mouse strains is attributed to loci other than *Tas1r3*. Taste responses to glycine provide a remarkable example: although there are substantial differences among mouse strains in responses to glycine [50,65], this variation is not attributed to the *Tas1r3* genotypes [31,69]. Consistent with the mouse work suggesting effects of genes other than *Tas1r3*, variation in saccharin preferences in rats is not associated with sequence variants of the rat *Tas1r3* gene [94], and therefore it must be attributed to the effects of other genes (see also [95]).

Because a sweet taste receptor appears to be a heterodimer of TIR2 and TIR3 proteins, sequence variants of the *Tas1r2* gene potentially can also result in variation in sweet taste responses. There is no evidence that *Tas1r2* variation contributes to mouse strain differences in responses to sweeteners. However, sequence variation among the *Tas1r2* orthologs underlies some species differences in sweet taste perception. Inability of rodents to perceive sweetness of several compounds that taste sweet to humans is attributed to variation between human and rodent orthologs of *Tas1r2* [87,96]. Lack of preference for sugars and some other sweeteners in cats (Felidae) [97,98] is due to pseudogenization of the *Tas1r2* gene, and thus inability to produce a TIR2 + TIR3 heterodimeric sweet taste receptor [8].

One of the genetic loci affecting sweet taste responses is *dpa* (D-phenylalanine aversion), which affects ability of mice to generalize a CTA between D-phenylalanine and sucrose, inferring that *dpa* affects ability to detect sweetness of D-phenylalanine. The *dpa* locus also affects responses of sucrose-sensitive fibers of the CT nerve to D-phenylalanine. B6 mice carry a dominant allele of *dpa* that determines an ability to recognize the sweetness of D-phenylalanine, whereas BALB/c mice carry a recessive *dpa* allele conferring inability to detect D-phenylalanine sweetness. The *dpa* locus was mapped to proximal Chr 4, a region distinct from the subtelomeric Chr 4 harboring the *Tas1r* genes [99–102]. It was suggested that the *dpa* locus can also affect responses to sweeteners in two-bottle tests [51]. Consistent with this, a locus on proximal Chr 4, in the *dpa* region, was found to be suggestively linked to consumption of, and CT responses to, sucrose [75]. An epistatic interaction between effects on sucrose intake of this locus and the *Tas1r3* locus suggests that these two loci may encode interacting components of sweet taste transduction [75].

In summary, these data show that sweetener preference has a complex genetic determination. In addition to the *Tas1r3* gene, other genetic loci play the role in genetic variation of taste responses to sweeteners. These other genes may encode additional peripheral taste transduction elements, including novel taste receptors. Alternatively, they may be involved in central mechanisms of ingestive behavioral responses or feedback mechanisms of modulation of sweet taste responses.

Umami taste genetics

Umami is a taste quality exemplified by taste of MSG and evoked also by some amino acids and purine 5'-nucleotides. There is a strong evidence that a heterodimer of T1R1 and T1R3 proteins functions as an umami taste receptor in humans and is a more broadly tuned in rodents to respond to L-amino acids (Table 2) [87,88,90,92]. A splice variant of a metabotropic glutamate receptor, mGluR4, was also proposed as a taste receptor for glutamate [103].

Little is known about genetic variation in umami taste responses. Humans appear to differ in perception of glutamate taste [104], but it is not known whether there is a genetic basis for this. A comparison of the CT nerve responses in three inbred mouse strains has shown differences in a synergistic effect between MSG and 5'-guanylate [105].

In the long-term two-bottle preference tests, mice from the B6 strain consumed more MSG and IMP than did mice from the 129 strain [106]. The strain difference in MSG consumption was in the opposite direction to the strain difference in NaCl consumption [66,107]. Although the B6 mice have higher avidity for both MSG and sweeteners than do the 129 mice [65,66,106], there is no correlation between preferences for these solutions in the F₂ hybrids derived from these two strains [106]. Thus, differences in MSG consumption between B6 and 129 mice are not related to the strain differences in salty or sweet taste responsiveness.

The role of the afferent gustatory input in these strain differences was examined by measuring integrated responses of the CT and IX nerves to umami taste stimuli. In the CT, responses to MSG and monoammonium L-glutamate were similar in B6 and 129 mice, but responses to IMP and guanosine-5'-monophosphate were lower in B6 than in 129 mice. Responses to umami stimuli in the IX nerve did not differ between the B6 and 129 strains [108]. Thus, the increased ingestive responses to the umami stimuli in B6 mice are accompanied by either unchanged or decreased neural responses to these stimuli. Lack of support for the role of the gustatory nerves in the enhanced consumption of MSG and IMP by B6 mice suggests that it is due to some other factors. A prior exposure to MSG affects subsequent MSG consumption [106], suggesting that it can be modulated by postingestive effects. The strain differences in gustatory neural responses to nucleotides but not glutamate [108] suggest that these compounds may activate distinct taste transduction mechanisms.

The T1R3 protein is involved in transduction of both sweet and umami tastes, and a disruption of the *Tas1r3*

gene diminishes behavioral or neural responses to umami taste stimuli [86-88,90]. Hence, variation of the *Tas1r3* gene might affect umami taste responses. However, an analysis of the F₂ hybrids between the B6 and 129 inbred mouse strains has shown that the *Tas1r3* allelic variants do not affect behavioral or neural taste responses to umami stimuli [69]. Thus, although the T1R3 receptor is involved in transduction of umami taste, the B6/129 sequence variants affecting its sensitivity to sweeteners do not affect its sensitivity to umami compounds.

Bitter taste

Bitter taste receptor genes

A large and diverse array of molecules evoke the sensation of bitterness, and the ability to detect and avoid these stimuli is assumed to have evolved as a mechanism to prevent ingestion of toxic foods [109]. Indeed, a strong correlation exists in the animal kingdom between bitter sensitivity and tolerance to toxic compounds [110]. A *Tas2r* family of G-protein coupled receptor genes linked to bitter taste sensitivity was identified in 2000 by laboratories led by Nicholas Ryba, Charles Zuker [111,112] and Linda Buck [113]. The T2R receptors are related to class A GPCRs and are characterized by a short N terminus with a potential transmembrane ligand-binding domain. To date, about 28 human and 36 mouse intact bitter taste receptor genes have been identified, although only a few have been de-orphanized (Table 2) [114].

The PTC polymorphism in humans and mice

The first example of a genetic basis to taste ability involved the serendipitous discovery of human differences in sensitivity to the bitter-tasting compound PTC [115]. Population surveys indicated a dimorphism in taste sensitivity for this substance, with individuals falling into two categories: "tasters" and "nontasters" [116,117]. Pedigree analysis suggested phenotypic control by a single, autosomal locus [118,119] although more recent studies postulated that PTC inheritance was better described by a polygenic model [120,121]. Linkage studies in humans supported associations with several chromosomal loci, including the KEL blood group antigen on Chr 7q (e.g. [122,123]). Dennis Drayna and colleagues [124,125] have shown that this locus corresponds to a polymorphic bitter taste receptor gene *TAS2R38* responsible for up to 85% of the variation in PTC sensitivity among human subjects. A secondary QTL on Chr 16p also has an effect on PTC aversion, although the nature of this gene is not known [124].

From the early discovery of human variation in PTC taste sensitivity, investigators proceeded to animal models of bitter taste genetics. Richter and Clisby [126] reported individual variation in PTC avoidance thresholds of rats, but the genetic basis for this was not pursued. Klein and

Figure 3

Aversion to PTC in taster mice develops over several days. Preference ratios (mean $\pm$ SE) for BALB/cBy (BALB), C3HeB/Fe (C3), and BALB $\times$ C3 F₁ (n = 10/strain) to 0.03 mM PTC over six consecutive 48-h tests. The dotted line represents a preference score of 0.5, which indicates equivalent consumption from solution and water tubes. Preference ratios for BALB and F₁ mice decreased following the initial test period; by the final test period these mice were strongly avoiding PTC. C3 mice remained indifferent to the taste of PTC across the entire test period. Strain differences in "developed aversion" to PTC have a genetic basis, with complete dominance of the avoider phenotype. However, the avoidance phenotype does not necessarily reflect increased bitter taste sensitivity. Figure from Boughter (unpublished).

DeFries [127] found that two-bottle intake of PTC varied in mice, with BALB/c mice significantly more sensitive to the bitter taste of PTC than other inbred strains, including C57BL/6J. A Mendelian cross (F₁, F₂, F₃, B₁, and B₂ generations) between these two strains yielded segregation ratios consistent with phenotypic control by a single autosomal locus, with a dominant taster allele. It was later recognized that PTC avoidance by BALB/c mice generally develops across 10–12 consecutive days of testing [128] (Figure 3). PTC is extremely toxic to mice (oral LD₅₀ 10 mg/kg), comparable to the rodent poison strychnine (oral LD₅₀ 2 mg/kg). Confining stimulus access to shorter durations helps to minimize or eliminate post-ingestive consequences (e.g. [29,129–132]). A direct comparison of intake tests and brief-access procedures demonstrated that differential avoidance of PTC among inbred strains depended on the amount of stimulus consumed, and not necessarily on immediate orosensory or taste cues [133]. When injected intraperitoneally, PTC was nearly as effective as LiCl in serving as the unconditioned stimulus in a CTA experiment [134]. Strain differences in PTC avoidance that develop over time therefore likely reflect other variables in addition to sensory ability, such as sensitivity

to its toxic effects, or differences in the speed of acquisition of a CTA.

Sucrose octaacetate (SOA) and quinine aversions and bitter taste receptor genes

The genetic basis of differences in intake of bitter compounds other than PTC has been studied throughout the last several decades, including substantial contributions from the laboratories of Ian Lush and Glayde Whitney (see [135,136]). Inbred strains differ in sensitivity to the commonly-used bitter stimuli, quinine and SOA, assessed using two-bottle preference tests [66,137–140]. For SOA, the strain difference is particularly robust as it occurs over a 2–3 log-step concentration range. SOA aversion appeared to be a fairly unambiguous model of monogenic control: Aversion ratios among segregating generations of mice were consistent with variation at a single autosomal locus (*Soa*, sucrose octaacetate aversion), with the taster allele dominant over either a "demitaster" (partial sensitivity) or nontaster allele [138,141,142]. Results of both brief-access behavioral tests and peripheral taste nerve electrophysiology yielded concentration-by-concentration similarity with two-bottle data, confirming that the

strain difference was gustatory in nature [132,140,143]. Relevant to this finding is the fact that SOA is a non-toxic, apparently non-odorous compound. *Soa* was subsequently mapped to distal mouse Chr 6 near proline-rich salivary protein genes *Prp2* and *Prh1* [136,144,145]. At least one other study has found evidence for polygenic control of SOA aversion [146], and interestingly, the aversion phenotype is subject to environmental modulation: unlike control tasters, taster mice raised on drinking water adulterated with a normally avoided concentration of SOA display only minimal avoidance of SOA in a two-bottle test with water [147].

Unlike responses to SOA, responses to quinine in two-bottle tests were demonstrated to be under polygenic control in segregating crosses or recombinant inbred strains [148-150]. However, linkage studies positioned a locus with major effects on quinine intake, dubbed *Qui* (quinine sensitivity, taste), in a close proximity with the *Prp2* and *Prh1* genes on distal Chr 6 [139,151,152]. Lush also reported linkage to this region for aversion to other bitter compounds, including cycloheximide, raffinose undecaacetate, and copper glycinate (Figure 4; [136,151,153]) and sagely hypothesized the existence of a cluster of bitter taste genes at this location, which "may have evolved from one original bitterness gene by a process of local duplication and differentiation" [136]. A recently characterized bitter taste receptor gene family in mice (36 *Tas2r* genes) forms two clusters on distal Chr 6 and one on Chr 15, including 24 genes at the *Soa* and *Qui* loci that collectively comprise a large haplotype for quinine taste sensitivity (Figure 5) [30,111,113,114]. Indeed, analysis suggests extensive gene duplication among the *Tas2r* genes [114]. Physical mapping of the *Soa-Qui* locus in strains of quinine taster (B6) and "non-taster" (DBA/2J) mice reveals considerable variance across the *Tas2r* genes [30]: Only two of 24 alleles were identical in both strains at the amino acid level. Furthermore, several of genes were either pseudogenes or deleted altogether in the nontaster strain, although a specific receptor-ligand relationship has not yet been uncovered.

Among bitter receptor genes in the mouse, only two *Tas2rs* have been de-orphanized thus far with respect to ligand, one responding exclusively to cycloheximide, the other to PROP and denatonium [5,154,155]. Several human T2R receptors, including TAS2R38, have also been characterized with respect to ligand binding (Table 2).

Evolution of the bitter taste receptor gene family

Prominent features of the bitter receptor gene family in vertebrates include a high rate of non-synonymous (i.e. resulting in amino acid change) substitutions, and a relatively conserved family size (~15 – 33 functional genes) with multiple orthologues among mammalian species

[156]. The most divergent portions of the *Tas2r* genes are those that code for extracellular domains, a possible site of ligand binding [114,157]. The particular nature of the high rate of mutations in this gene family has led to speculation of relaxation of selective constraint and loss of function in primate species [158,159]. On the other hand, analysis of TAS2R haplotypes among humans shows a greater diversity than expected, and suggests strong effects of natural selection [6,160]. The number of functional receptors may have decreased in humans and other primates as the chemosensory function of these species has diminished in importance relative to other mammals, but positive selection has shaped the evolution of bitter taste receptors in response to variation in local plant species that are toxic.

Salt taste

A prototypical salty taste stimulus is NaCl, but several other compounds (e.g., chlorides and other salts of Na, Li and K) also evoke salty taste. Sodium enters epithelial cells through the sodium-selective epithelial channel ENaC. ENaC is a member of the degenerin/ENaC superfamily of ion channels and is inhibited by a diuretic, amiloride. Taste receptor cells express amiloride-blockable, Na⁺-selective channels that share a number of physiological properties with the ENaCs (for review, see [161]). Thus, a large portion of NaCl taste is likely dependent on the influx of sodium through ENaC. However, whether ENaC plays the role of a salt taste receptor requires further studies.

Sodium is an important nutrient, and its consumption is under a strong homeostatic control [162,163]. Therefore, results of the long-term preference tests with sodium salts may be affected not only by perception of their taste, but also by postingestive effects. Thus, additional evidence is usually needed to link variation in salt consumption with differences in salty taste perception.

Salt consumption may be under genetic control in humans [164]. Inbred strain differences in voluntary NaCl consumption have been reported in both mice [28,66,107,166-169] and rats [170-174]. Although these genetic differences in NaCl intake or preference are often profound, it is not clear whether they originate from differences in taste perception or are due to other factors. A few studies undertook a genetic analyses of NaCl consumption in mice [73] and rats [175-179]. In rats, salt intake was found to be linked to the Y Chr [177,180]. No linkage data for NaCl consumption in mice have been published.

Despite large strain variation in NaCl preferences, most studies found either small or no differences in integrated responses of the gustatory nerves to NaCl among strains of

Figure 4

Genome-wide interval mapping of taste responses to bitter and sweet compounds: 0.1 – 0.4 mg/ml quinine, 0.3 mM beta-lactose acetate (BLA), 0.4 mM raffinose undecaacetate (RUA), 10 mM copper glycinate and 3.2 mM saccharin. LRS = Likelihood Ratio Score. The taste stimuli were tested in the two-bottle preference tests in the BXD recombinant inbred mouse strains. Phenotype data were previously published [46, 72, 151, 153] and are available at <http://www.genenetwork.org>. Mapping was conducted using WebQTL software <http://www.genenetwork.org>. The interval mapping illustrates previously published linkages for avoidance of all four bitter-tasting stimuli to Chr 6 and for saccharin preference to Chr 4 (*Sac/Tas1r3* locus). A recent study of bitter taste in BXD mice using brief-access procedures demonstrates that the QTL on Chr 6 reflects orosensory processing, and not post-ingestive effects, of the bitter stimulus quinine [30].

mice [108,166,181-183] and rats [184,185], with some exceptions [186].

In rodents, lingual application of amiloride suppresses the chorda tympani responses to NaCl [187]. There are mouse and rat strain differences in sensitivity of the chorda tympani response to amiloride. Initial studies suggested that amiloride sensitivity of the CT nerve response to NaCl is associated with behavioral responses to NaCl: strains with stronger amiloride-induced suppression of the neural response to NaCl had stronger NaCl avoidance [166,182,183,186]. However, this relationship was not found in all studies. For example, NZB/BlNJ and CBA/J mice dramatically differ in NaCl consumption [107,168] but have chorda tympani responses to NaCl of similar magnitude and sensitivity to amiloride [181]. Lack of correspondence between amiloride sensitivity of the CT response to NaCl and NaCl consumption was also found when three strains of rats, Fisher 344, Wistar and Sprague-Dawley, were compared [185,186]. Furthermore, amiloride was found to affect NaCl perception in mice regardless of amiloride sensitivity of their gustatory neural responses to NaCl [32,188]. More studies are needed to determine whether there is a mechanistic relationship between amiloride sensitivity of neural responses to NaCl and NaCl preference. It appears that the genetic variation in voluntary NaCl consumption can depend on a variety

of factors, including, but not limited to, peripheral taste responsiveness.

Rats typically prefer NaCl solutions of intermediate (osmotically hypotonic or isotonic) concentrations. However, rats from the Fisher 344 strain do not display NaCl preferences [170]. A series of subsequent studies have presented a detailed analysis of mechanisms underlying this lack of NaCl preference in Fisher 344 rats. Relative to rats with typical responses to NaCl, Fisher 344 rats develop attenuated salt appetite in response to sodium deficiency or to interference with renin-angiotensin-aldosterone system, but they have similar sodium loss during sodium depletion. This suggests that differences in hedonic perception of NaCl rather than differences in sodium metabolism underlie aberrant NaCl consumption by the Fisher 344 rats [189-191]. Consistent with this, Fisher 344 rats show decreased appetitive and increased aversive oral motor responses to NaCl [192], and their NaCl aversion is abolished by CT, but not IX nerve transection [193,194]. However, NaCl detection thresholds were not altered in Fisher 344 rats [195]. Thus, it appears that NaCl aversion in Fisher 344 rats is mediated by aversive gustatory information conveyed by the CT nerve.

Increased NaCl consumption by SHR (spontaneously hypertensive) rats does not depend on strain differences in peripheral gustatory input [184,196], but appears to be

Figure 5
A map of the cluster of the *Tas2r* bitter taste genes on distal mouse Chr 6. Twenty-four intact *Tas2r* genes map to distal Chr 6 (black). The *Tas2rs* are found in two sub-clusters on either side of the polymorphic marker *D6Mit13* (red) and two genes encoding proline-rich salivary proteins (*Prp2* and *Prh1*; red). Map positions are in Mb (Build 33 assembly of the B6 genome). Figure copyright [30].

mediated by the brain renin-angiotensin system [197,198].

Taste and other complex phenotypes

Taste perception involves hedonic processes and is an important factor affecting ingestive behavior. Therefore, genetic variation in taste preferences is likely to affect complex phenotypes that depend on oral consumption of nutrients or drugs, or involve pleasure-seeking behavior. There are several examples of association of taste and complex traits in humans, including relationship between sweet taste and obesity [34,41,199,200] or alcohol intake [201-205], and between bitter taste and alcohol intake [206-212], smoking [213,214], food choice [215] and other health-related traits [216]. Studies of rodents elucidated some genetic factors and physiological mechanisms for association between sweet taste and alcohol, and have potential for unveiling mechanisms of the other associations between taste and complex phenotypes.

The oral consumption of alcohol is accompanied by chemosensory perception of its flavor, which plays an important role in its acceptance and rejection. Three independent sensory systems – taste, olfaction and chemosensory irritation – are involved in the perception of alcohol flavor. Humans perceive alcohol as a combination of sweet and bitter tastes, odors and oral irritation (e.g., burning sensation), all of which vary as a function of concentration [209,217,218]. Likewise, rodents detect the sweet (sucrose-like) and bitter (quinine-like) taste [219-221], odor volatiles [222,223] of alcohol, and probably the other components detected by humans [224,225].

Perception of the sweet taste component of ethanol by rodents was shown in behavioral and neurophysiological experiments. CTAs generalize between ethanol and sucrose [219,221,223,226,227]. Electrophysiological recordings indicate that lingual application of ethanol activates sweet-best neural fibers in the gustatory nerves [228,229] and sweet-best units in the nucleus of the tractus solitarius [230,231]; this activity is blocked by application of gumarin, a peripheral antagonist of sweet taste [231]. Central mechanisms that determine hedonic responses to ethanol and sweeteners also overlap and involve opioidergic, serotonergic and dopaminergic brain neurotransmitter systems [232-236]. In addition, there may be common signals related to the caloric value of ethanol and sugars [237-242].

Positive correlations between preferences for ethanol and sweeteners in rats and mice were found among various strains and in segregating crosses [25,45,52,66,72,73,243-246], reviewed in [62,63]. This genetically determined association can be underlaid by any of the mechanisms

described above, including peripheral or central taste processing, or postingestive reward.

Genetic analysis of a cross between mice from a high ethanol- and sweetener-preferring B6 strain and a low ethanol- and sweetener-preferring 129 strain suggested that the strain differences in sweetener and ethanol consumption depend on relatively small and partially overlapping sets of genes [73]. One of these genetic loci, *Ap3q* (alcohol preference 3 QTL), maps to the subtelomeric region of Chr 4 [93] overlapping with the saccharin preference (*Sac*) locus that corresponds to the sweet taste receptor gene, *Tas1r3* [77]. This suggests that the *Tas1r3* gene is identical to the *Ap3q* locus and that its pleiotropic effect on ethanol consumption is mediated by genetic differences in perception of the sweet taste component of ethanol flavor: higher hedonic attractiveness of ethanol sweetness results in higher ethanol intake by B6 mice. The role of the T1R3 receptor in alcohol consumption was confirmed in a recent study showing that mutant mice lacking the *Tas1r3* gene have diminished ethanol intakes and preferences [247]. In addition to the *Tas1r3* gene, there are other genetic loci with pleiotropic effects on ethanol and sweetener intake [248,249].

Conclusion

Behavioral genetic studies of taste responsiveness have been instrumental in the discovery of G-protein-coupled taste receptors. Genetic mapping of a human PTC/PROP taste sensitivity locus and mouse bitter and sweet taste loci have facilitated identification of the T1R and T2R families of taste receptors. The behavioral genetic approach has potential for more discoveries of taste mechanisms. There is strong evidence that multiple genes control behavioral taste responses. These yet unknown genes are likely to be involved in different stages of the taste processing pathway, including taste reception, transduction and transmission in the periphery and in the brain, and interaction of taste processing with homeostatic systems involved in the regulation of feeding, appetite, fluid balance, and reward. Genetics has experienced dramatic progress in recent years, with genome sequencing completed for several species, including mouse and human. These advances in genomic resources tremendously facilitate chromosomal mapping of genes affecting taste responsiveness and their identification. This turns behavioral genetics into a powerful approach for understanding mechanisms of taste.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JDB and AAB contributed equally to this review.

Acknowledgements

Supported by NIH grants DC00882 (G.K. Beauchamp), AA11028 (AAB), DC000353 and NS052366 (JDB), and an Ajinomoto Amino Acid Research Program Focused Research grant (AAB).

This article has been published as part of *BMC Neuroscience* Volume 8 Supplement 3, 2007: The chemical senses: recent advances and new promises. The full contents of the supplement are available online at <http://www.biomedcentral.com/1471-2202/8?issue=S3>.

References

1. Bachmanov AA, Beauchamp GK: **Taste Receptor Genes.** *Annu Rev Nutr* 2007, **27**:389-414.
2. Drayna D: **Human taste genetics.** *Annu Rev Genomics Hum Genet* 2005, **6**:217-235.
3. Mombaerts P: **Genes and ligands for odorant, vomeronasal and taste receptors.** *Nat Rev Neurosci* 2004, **5**:263-278.
4. Scott K: **Taste recognition: food for thought.** *Neuron* 2005, **48**:455-464.
5. Mueller KL, Hoon MA, Erlenbach I, Chandrashekar J, Zuker CS, Ryba NJ: **The receptors and coding logic for bitter taste.** *Nature* 2005, **434**:225-229.
6. Kim UK, Drayna D: **Genetics of individual differences in bitter taste perception: lessons from the PTC gene.** *Clin Genet* 2005, **67**:275-280.
7. Maruyama Y, Pereira E, Margolskee RF, Chaudhari N, Roper SD: **Umami responses in mouse taste cells indicate more than one receptor.** *J Neurosci* 2006, **26**:2227-2234.
8. Li X, Li W, Wang H, Cao J, Maehashi K, Huang L, Bachmanov AA, Reed DR, Legrand-Defretin V, Beauchamp GK, Brand JG: **Pseudogenization of a sweet-receptor gene accounts for cats' indifference toward sugar.** *PLoS Genet* 2005, **1**:27-35.
9. Scott TR: **The role of taste in feeding.** *Appetite* 2001, **37**:111-113.
10. Cho YK, Li CS, Smith DV: **Descending influences from the lateral hypothalamus and amygdala converge onto medullary taste neurons.** *Chem Senses* 2003, **28**:155-171.
11. Hajnal A, Takenouchi K, Norgren R: **Effect of intraduodenal lipid on parabrachial gustatory coding in awake rats.** *J Neurosci* 1999, **19**:7182-7190.
12. Atchley DP, Weaver KL, Eckel LA: **Taste responses to dilute sucrose solutions are modulated by stage of the estrous cycle and fenfluramine treatment in female rats.** *Physiol Behav* 2005, **86**:265-271.
13. Curtis KS, Stratford JM, Contreras RJ: **Estrogen increases the taste threshold for sucrose in rats.** *Physiol Behav* 2005, **86**:281-286.
14. Simon SA, Liu L, Erickson RP: **Neuropeptides modulate rat chorda tympani responses.** *Am J Physiol Regul Integr Comp Physiol* 2003, **284**:R1494-R1505.
15. Shigemura N, Ohta R, Kusakabe Y, Miura H, Hino A, Koyano K, Nakashima K, Ninomiya Y: **Leptin modulates behavioral responses to sweet substances by influencing peripheral taste structures.** *Endocrinology* 2004, **145**:839-847.
16. Mook DG: **Oral and postingestional determinants of the intake of various solutions in rats with esophageal fistulas.** *Journal of Comparative and Physiological Psychology* 1963, **56**:645-659.
17. Marella S, Fischler W, Kong P, Asgarian S, Rueckert E, Scott K: **Imaging taste responses in the fly brain reveals a functional map of taste category and behavior.** *Neuron* 2006, **49**:285-295.
18. Scott K, Brady R Jr, Cravchik A, Morozov P, Rzhetsky A, Zuker C, Axel R: **A chemosensory gene family encoding candidate gustatory and olfactory receptors in Drosophila.** *Cell* 2001, **104**:661-673.
19. Huang AL, Chen X, Hoon MA, Chandrashekar J, Guo W, Trankner D, Ryba NJ, Zuker CS: **The cells and logic for mammalian sour taste detection.** *Nature* 2006, **442**:934-938.
20. Ishimaru Y, Inada H, Kubota M, Zhuang H, Tominaga M, Matsunami H: **Transient receptor potential family members PKD1L3 and PKD2L1 form a candidate sour taste receptor.** *Proc Natl Acad Sci U S A* 2006, **103**:12569-12574.
21. Harris H, Kalmus H: **The measurement of taste sensitivity to phenylthiourea (P.T.C.).** *Annals of Eugenics* 1949, **15**:24-31.
22. Bartoshuk LM, Duffy VB, Miller IJ: **PTC/PROP tasting: anatomy, psychophysics, and sex effects.** *Physiol Behav* 1994, **56**:1165-1171.

23. Hansen JL, Reed DR, Wright MJ, Martin NG, Breslin PA: **Heritability and Genetic Covariation of Sensitivity to PROP, SOA, Quinine HCl, and Caffeine.** *Chem Senses* 2006, **31**:403-413.
24. Spector AC: **Psychophysical evaluation of taste function in nonhuman mammals.** In *Handbook of Olfaction and Gustation* Edited by: Doty RL. New York: Marcel Dekker; 2003:869-879.
25. McClearn GE, Rodgers DA: **Differences in alcohol preference among inbred strains of mice.** *Quart J Stud Alcohol* 1959, **20**:691-695.
26. Nachman M: **The inheritance of saccharin preference.** *J Comp Physiol Psychol* 1959, **52**:451-457.
27. Hoshishima K, Yokoyama S, Seto K: **Taste sensitivity in various strains of mice.** *Am J Physiol* 1962, **202**:1200-1204.
28. Lush IE: **The genetics of bitterness, sweetness, and saltiness in strains of mice.** In *Genetics of Perception and Communication Volume 3*. Edited by: Wysocki CJ, Kare MR. New York: Marcel Dekker; 1991:227-241. *Chemical Senses*
29. Glendinning JJ, Gresack J, Spector AC: **A high-throughput screening procedure for identifying mice with aberrant taste and oromotor function.** *Chem Senses* 2002, **27**:461-474.
30. Nelson TM, Munger SD, Boughter JD Jr: **Haplotypes at the Tas2r locus on distal chromosome 6 vary with quinine taste sensitivity in inbred mice.** *BMC Genet* 2005, **6**:32.
31. Eylam S, Spector AC: **Stimulus processing of glycine is dissociable from that of sucrose and glucose based on behaviorally measured taste signal detection in Sac 'taster' and 'non-taster' mice.** *Chem Senses* 2004, **29**:639-649.
32. Eylam S, Spector AC: **Taste discrimination between NaCl and KCl is disrupted by amiloride in inbred mice with amiloride-insensitive chorda tympani nerves.** *Am J Physiol Regul Integr Comp Physiol* 2005, **288**:R1361-1368.
33. Gent JF, Bartoshuk LM: **Sweetness of sucrose, neohesperidin dihydrochalcone, and saccharin is related to genetic ability to taste the bitter substance 6-n-propylthiouracil.** *Chemical Senses* 1983, **7**:265-272.
34. Reed DR, Bachmanov AA, Beauchamp GK, Tordoff MG, Price RA: **Heritable variation in food preferences and their contribution to obesity.** *Behavior Genetics* 1997, **27**:373-387.
35. Looy H, Weingarten HP: **Facial expressions and genetic sensitivity to 6-n-propylthiouracil predict hedonic response to sweet.** *Physiol Behav* 1992, **52**:75-82.
36. Looy H, Callaghan S, Weingarten HP: **Hedonic response of sucrose likers and dislikers to other gustatory stimuli.** *Physiol Behav* 1992, **52**:219-225.
37. Beauchamp GK, Moran M: **Dietary experience and sweet taste preference in human infants.** *Appetite* 1982, **3**:139-152.
38. Desor JA, Greene LS, Maller O: **Preferences for sweet and salty in 9- to 15-year-old and adult humans.** *Science* 1975, **190**:686-687.
39. Greene LS, Desor JA, Maller O: **Heredity and experience: their relative importance in the development of taste preference in man.** *J Comp Physiol Psychol* 1975, **89**:279-284.
40. Falciglia GA, Norton PA: **Evidence for a genetic influence on preference for some foods.** *J Am Diet Assoc* 1994, **94**:154-158.
41. McDaniel AH, Reed DR: **The human sweet tooth and its relationship to obesity.** In *Genomics and Proteomics in Nutrition* Edited by: Berdanier CD, Moustaid-Moussa N. New York: Marcel Dekker, Inc; 2003:51-70.
42. Reed DR, Li X, Bachmanov AA, Mascioli K, Beauchamp GK: **The molecular basis of the mammalian sweet tooth.** In *Progress in Obesity Research Volume 9*. Ltd. JLE. London; 2003:304-306. G Medeiros-Neto AH, and C. Bouchard (Series Editor)
43. Lewis SR, Ahmed S, Dym C, Khaimova E, Kest B, Bodnar RJ: **Inbred mouse strain survey of sucrose intake.** *Physiol Behav* 2005, **85**:546-556.
44. Fuller JL: **Single-locus control of saccharin preference in mice.** *J Hered* 1974, **65**:33-36.
45. Rodgers DA, McClearn GE: **Sucrose versus ethanol appetite in inbred strains of mice.** *Q J Stud Alcohol* 1964, **25**:26-35.
46. Lush IE: **The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose.** *Genetical Research* 1989, **53**:95-99.
47. Ramirez I, Fuller JL: **Genetic influence on water and sweetened water consumption in mice.** *Physiol Behav* 1976, **16**:163-168.
48. Pelz WE, Whitney G, Smith JC: **Genetic influences on saccharin preference of mice.** *Physiol Behav* 1973, **10**:263-265.
49. Stockton MD, Whitney G: **Effects of genotype, sugar, and concentration on sugar preference of laboratory mice (Mus musculus).** *J Comp Physiol Psychol* 1974, **86**:62-68.
50. Lush IE, Hornigold N, King P, Stoye JP: **The genetics of tasting in mice. VII. Glycine revisited, and the chromosomal location of Sac and Soa.** *Genetical Research* 1995, **66**:167-174.
51. Capeless CG, Whitney G: **The genetic basis of preference for sweet substances among inbred strains of mice: Preference ratio phenotypes and the alleles of the Sac and dpa loci.** *Chemical Senses* 1995, **20**:291-298.
52. Belknap JK, Crabbe JC, Young ER: **Voluntary consumption of alcohol in 15 inbred mouse strains.** *Psychopharmacology* 1993, **112**:503-510.
53. Reed DR, Li S, Li X, Huang L, Tordoff MG, Starling-Roney R, Taniguchi K, West DB, Ohmen JD, Beauchamp GK, Bachmanov AA: **Polymorphisms in the taste receptor gene (Tas1r3) region are associated with saccharin preference in 30 mouse strains.** *J Neurosci* 2004, **24**:938-946.
54. Kotlus BS, Blizard DA: **Measuring gustatory variation in mice: A short-term fluid-intake test.** *Physiol Behav* 1998, **64**:37-47.
55. Dotson CD, Spector AC: **The relative affective potency of glycine, L-serine and sucrose as assessed by a brief-access taste test in inbred strains of mice.** *Chem Senses* 2004, **29**:489-498.
56. Glendinning JJ, Chyou S, Lin I, Onishi M, Patel P, Zheng KH: **Initial licking responses of mice to sweeteners: effects of tas1r3 polymorphisms.** *Chem Senses* 2005, **30**:601-614.
57. Ninomiya Y, Higashi T, Katsukawa H, Mizukoshi T, Funakoshi M: **Qualitative discrimination of gustatory stimuli in three different strains of mice.** *Brain Research* 1984, **322**:83-92.
58. Inoue M, McCaughey SA, Bachmanov AA, Beauchamp GK: **Whole-nerve chorda tympani responses to sweeteners in C57BL/6ByJ and 129P3/J mice.** *Chemical Senses* 2001, **26**:915-923.
59. Frank ME, Blizard DA: **Chorda tympani responses in two inbred strains of mice with different taste preferences.** *Physiol Behav* 1999, **67**:287-297.
60. Ninomiya Y, Mizukoshi T, Higashi T, Katsukawa H, Funakoshi M: **Gustatory neural responses in three different strains of mice.** *Brain Research* 1984, **302**:305-314.
61. Dess NK, Minor TR: **Taste and emotionality in rats selectively bred for high versus low saccharin intake.** *Animal Learning and Behavior* 1996, **24**:105-115.
62. Overstreet DH, Rezvani AH, Parsian A: **Behavioural features of alcohol-preferring rats: focus on inbred strains.** *Alcohol Alcohol* 1999, **34**:378-385.
63. Kampov-Polevoy AB, Garbutt JC, Janowsky DS: **Association between preference for sweets and excessive alcohol intake: a review of animal and human studies.** *Alcohol Alcohol* 1999, **34**:386-395.
64. Frank ME, Wada Y, Makino J, Mizutani M, Umezawa H, Katsue Y, Hettinger TP, Blizard DA: **Variation in intake of sweet and bitter solutions by inbred strains of golden hamsters.** *Behav Genet* 2004, **34**:465-476.
65. Bachmanov AA, Tordoff MG, Beauchamp GK: **Sweetener preference in C57BL/6ByJ and 129P3/J mice.** *Chemical Senses* 2001, **26**:905-913.
66. Bachmanov AA, Tordoff MG, Beauchamp GK: **Ethanol consumption and taste preferences in C57BL/6ByJ and 129/J mice.** *Alcohol Clin Exp Res* 1996, **20**:201-206.
67. Bachmanov AA, Reed DR, Tordoff MG, Price RA, Beauchamp GK: **Nutrient preference and diet-induced adiposity in C57BL/6ByJ and 129P3/J mice.** *Physiol Behav* 2001, **72**:603-613.
68. Manita S, Bachmanov AA, Li X, Beauchamp GK, Inoue M: **Is glycine "sweet" to mice? Mouse strain differences in perception of glycine taste.** *Chem Senses* 2006, **31**:785-793.
69. Inoue M, Reed DR, Li X, Tordoff MG, Beauchamp GK, Bachmanov AA: **Allelic variation of the Tas1r3 taste receptor gene selectively affects behavioral and neural taste responses to sweeteners in the F₂ hybrids between C57BL/6ByJ and 129P3/J mice.** *J Neurosci* 2004, **24**:2296-2303.
70. Fuller JL: **Single-locus control of saccharin preference in mice.** *Journal of Heredity* 1974, **65**:33-36.
71. Belknap JK, Crabbe JC, Plomin R, McClearn GE, Sampson KE, O'Toole LA, Gora-Maslak G: **Single-locus control of saccharin intake in BXD/Ty recombinant inbred (RI) mice: Some methodological implications for RI strain analysis.** *Behavior Genetics* 1992, **22**:81-100.

72. Phillips TJ, Crabbe JC, Metten P, Belknap JK: **Localization of genes affecting alcohol drinking in mice.** *Alcohol Clin Exp Res* 1994, **18**:931-941.
73. Bachmanov AA, Reed DR, Tordoff MG, Price RA, Beauchamp GK: **Intake of ethanol, sodium chloride, sucrose, citric acid, and quinine hydrochloride solutions by mice: a genetic analysis.** *Behavior Genetics* 1996, **26**:563-573.
74. Blizard DA, Kotlus B, Frank ME: **Quantitative trait loci associated with short-term intake of sucrose, saccharin and quinine solutions in laboratory mice.** *Chemical Senses* 1999, **24**:373-385.
75. Bachmanov AA, Reed DR, Ninomiya Y, Inoue M, Tordoff MG, Price RA, Beauchamp GK: **Sucrose consumption in mice: major influence of two genetic loci affecting peripheral sensory responses.** *Mammalian Genome* 1997, **8**:545-548.
76. Li X, Inoue M, Reed DR, Huque T, Puchalski RB, Tordoff MG, Ninomiya Y, Beauchamp GK, Bachmanov AA: **High-resolution genetic mapping of the saccharin preference locus (Sac) and the putative sweet taste receptor (TIR1) gene (Gpr70) to mouse distal Chromosome 4.** *Mammalian Genome* 2001, **12**:13-16.
77. Bachmanov AA, Li X, Reed DR, Ohmen JD, Li S, Chen Z, Tordoff MG, de Jong PJ, Wu C, West DB, et al.: **Positional cloning of the mouse saccharin preference (Sac) locus.** *Chemical Senses* 2001, **26**:925-933.
78. Li X, Bachmanov AA, Li S, Chen Z, Tordoff MG, Beauchamp GK, de Jong PJ, Wu C, Chen L, West DB, et al.: **Genetic, physical and comparative map of the subtelomeric region of mouse chromosome 4.** *Mammalian Genome* 2002, **13**:5-19.
79. Wong GT, Gannon KS, Margolskee RF: **Transduction of bitter and sweet taste by gustducin.** *Nature* 1996, **381**:796-800.
80. Montmayeur JP, Liberles SD, Matsunami H, Buck LB: **A candidate taste receptor gene near a sweet taste locus.** *Nat Neurosci* 2001, **4**:492-498.
81. Kitagawa M, Kusakabe Y, Miura H, Ninomiya Y, Hino A: **Molecular genetic identification of a candidate receptor gene for sweet taste.** *Biochem Biophys Res Commun* 2001, **283**:236-242.
82. Hoon MA, Adler E, Lindemeier J, Battey JF, Ryba NJ, Zuker CS: **Putative mammalian taste receptors: a class of taste-specific GPCRs with distinct topographic selectivity.** *Cell* 1999, **96**:541-551.
83. Max M, Shanker YG, Huang L, Rong M, Liu Z, Campagne F, Weinstein H, Damak S, Margolskee RF: **Tas1r3, encoding a new candidate taste receptor, is allelic to the sweet responsiveness locus Sac.** *Nat Genet* 2001, **28**:58-63.
84. Sainz E, Korley JN, Battey JF, Sullivan SL: **Identification of a novel member of the TIR family of putative taste receptors.** *J Neurochem* 2001, **77**:896-903.
85. Nelson G, Hoon MA, Chandrashekar J, Zhang Y, Ryba NJ, Zuker CS: **Mammalian sweet taste receptors.** *Cell* 2001, **106**:381-390.
86. Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E: **Human receptors for sweet and umami taste.** *Proc Natl Acad Sci U S A* 2002, **99**:4692-4696.
87. Nelson G, Chandrashekar J, Hoon MA, Feng L, Zhao G, Ryba NJ, Zuker CS: **An amino-acid taste receptor.** *Nature* 2002, **416**:199-202.
88. Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ, Zuker CS: **The receptors for mammalian sweet and umami taste.** *Cell* 2003, **115**:255-266.
89. Liao J, Schultz PG: **Three sweet receptor genes are clustered in human chromosome 1.** *Mamm Genome* 2003, **14**:291-301.
90. Damak S, Rong M, Yasumatsu K, Kokrashvili Z, Varadarajan V, Zou S, Jiang P, Ninomiya Y, Margolskee RF: **Detection of sweet and umami taste in the absence of taste receptor TIR3.** *Science* 2003, **301**:850-853.
91. Nie Y, Vignes S, Hobbs JR, Conn GL, Munger SD: **Distinct contributions of TIR2 and TIR3 taste receptor subunits to the detection of sweet stimuli.** *Curr Biol* 2005, **15**:1948-1952.
92. Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E: **Human receptors for sweet and umami taste.** *Proc Natl Acad Sci U S A* 2002, **99**:4692-4696.
93. Bachmanov AA, Reed DR, Li X, Li S, Beauchamp GK, Tordoff MG: **Voluntary ethanol consumption by mice: genome-wide analysis of quantitative trait loci and their interactions in a C57BL/6ByJ x 129P3/J F2 intercross.** *Genome Res* 2002, **12**:1257-1268.
94. Lu K, McDaniel AH, Tordoff MG, Li X, Beauchamp GK, Bachmanov AA, VanderWeele DA, Chapman CD, Dess NK, Huang L, et al.: **No relationship between sequence variation in protein coding regions of the Tas1r3 gene and saccharin preference in rats.** *Chem Senses* 2005, **30**:231-240.
95. Foroud T, Bice P, Castelluccio P, Bo R, Ritchotte A, Stewart R, Lumeng L, Li TK, Carr L: **Mapping of QTL influencing saccharin consumption in the selectively bred alcohol-preferring and -nonpreferring rat lines.** *Behav Genet* 2002, **32**:57-67.
96. Xu H, Staszewski L, Tang H, Adler E, Zoller M, Li X: **Different functional roles of TIR subunits in the heteromeric taste receptors.** *Proc Natl Acad Sci U S A* 2004, **101**:14258-14263.
97. Beauchamp GK, Maller O, Rogers JG: **Flavor preferences in cats (Felis catus and Panthera sp.).** *Journal of Comparative and Physiological Psychology* 1977, **91**:1118-1127.
98. Bartoshuk LM, Jacobs HL, Nichols TL, Hoff LA, Ryckman JJ: **Taste rejection of nonnutritive sweeteners in cats.** *J Comp Physiol Psychol* 1975, **89**:971-975.
99. Ninomiya Y, Sako N, Katsukawa H, Funakoshi M: **Taste receptor mechanisms influenced by a gene on chromosome 4 in mice.** In *Genetics of Perception and Communication Volume 3*. Edited by: Wysocki CJ, Kare MR. New York: Marcel Dekker; 1991:267-278. *Chemical Senses*
100. Ninomiya Y, Higashi T, Mizukoshi T, Funakoshi M: **Genetics of the ability to perceive sweetness of d-phenylalanine in mice.** *Annals of the New York Academy of Sciences* 1987, **510**:527-529.
101. Ninomiya Y, Nomura T, Katsukawa H: **Genetically variable taste sensitivity to D-amino acids in mice.** *Brain Res* 1992, **596**:349-352.
102. Shigemura N, Yasumatsu K, Yoshida R, Sako N, Katsukawa H, Nakashima K, Imoto T, Ninomiya Y: **The Role of the dpa Locus in Mice.** *Chem Senses* 2005, **30**(Suppl 1):i84-i85.
103. Chaudhari N, Landin AM, Roper SD: **A metabotropic glutamate receptor variant functions as a taste receptor.** *Nat Neurosci* 2000, **3**:113-119.
104. Lugaz O, Pillias AM, Faurion A: **A new specific ageusia: some humans cannot taste L-glutamate.** *Chem Senses* 2002, **27**:105-115.
105. Ninomiya Y, Kurenuma S, Nomura T, Uebayashi H, Kawamura H: **Taste synergism between monosodium glutamate and 5'-ribonucleotide in mice.** *Comp Biochem Physiol A* 1992, **101**:97-102.
106. Bachmanov AA, Tordoff MG, Beauchamp GK: **Intake of umami-tasting solutions by mice: a genetic analysis.** *Journal of Nutrition* 2000, **130**:935S-941S.
107. Bachmanov AA, Tordoff MG, Beauchamp GK: **Voluntary sodium chloride consumption by mice: Differences among five inbred strains.** *Behavior Genetics* 1998, **28**:117-124.
108. Inoue M, Beauchamp GK, Bachmanov AA: **Gustatory Neural Responses to Umami Taste Stimuli in C57BL/6ByJ and 129P3/J Mice.** *Chem Senses* 2004, **29**:789-795.
109. Spielman AI, Huque T, Whitney G, Brand JG: **The diversity of bitter taste signal transduction mechanisms.** In *Sensory Transduction* Edited by: Corey DP, Roper SD. New York: Rockefeller Univ. Press; 1992:307-324.
110. Glendinning JL: **Is the bitter rejection response always adaptive?** *Physiol Behav* 1994, **56**:1217-1227.
111. Adler E, Hoon MA, Mueller KL, Chandrashekar J, Ryba NJ, Zuker CS: **A novel family of mammalian taste receptors.** *Cell* 2000, **100**:693-702.
112. Chandrashekar J, Mueller KL, Hoon MA, Adler E, Feng L, Guo W, Zuker CS, Ryba NJ: **T2Rs function as bitter taste receptors.** *Cell* 2000, **100**:703-711.
113. Matsunami H, Montmayeur JP, Buck LB: **A family of candidate taste receptors in human and mouse.** *Nature* 2000, **404**:601-604.
114. Shi P, Zhang J, Yang H, Zhang YP: **Adaptive diversification of bitter taste receptor genes in Mammalian evolution.** *Mol Biol Evol* 2003, **20**:805-814.
115. Fox AL: **Six in ten "tasteblind" to bitter chemicals.** *Sci News Lett* 1931, **9**:249.
116. Fox AL: **The relationship between chemical constitution and taste.** *Proc Natl Acad Sci U S A* 1932, **18**:115-120.
117. Harris H, Kalmus H: **The measurement of taste sensitivity to phenylthiourea (P.T.C.).** *Annals of Eugenics* 1949, **15**:24-31.
118. Snyder LH: **Inherited taste deficiency.** *Science* 1931, **74**:151-152.

119. Blakeslee AF: **Genetics of sensory thresholds: Taste for phenylthio carbamide.** *Proc Natl Acad Sci U S A* 1932, **18**:120-130.
120. Olson JM, Boehnke M, Neiswanger K, Roche AF, Siervogel RM: **Alternative genetic models for the inheritance of the phenylthiocarbamide taste deficiency.** *Genet Epidemiol* 1989, **6**:423-434.
121. Reddy BM, Rao DC: **Phenylthiocarbamide taste sensitivity revisited: complete sorting test supports residual family resemblance.** *Genet Epidemiol* 1989, **6**:413-421.
122. Conneally PM, Dumont-Driscoll M, Huntzinger RS, Nance WE, Jackson CE: **Linkage relations of the loci for Kell and phenylthiocarbamide taste sensitivity.** *Hum Hered* 1976, **26**:267-271.
123. Reed DR, Nanthakumar E, North M, Bell C, Bartoshuk LM, Price RA: **Localization of a gene for bitter-taste perception to human chromosome 5p15.** *Am J Hum Genet* 1999, **64**:1478-1480.
124. Drayna D, Coon H, Kim UK, Elsner T, Cromer K, Otterud B, Baird L, Peiffer AP, Leppert M: **Genetic analysis of a complex trait in the Utah Genetic Reference Project: a major locus for PTC taste ability on chromosome 7q and a secondary locus on chromosome 16p.** *Hum Genet* 2003, **112**:567-572.
125. Kim UK, Jorgenson E, Coon H, Leppert M, Risch N, Drayna D: **Positional cloning of the human quantitative trait locus underlying taste sensitivity to phenylthiocarbamide.** *Science* 2003, **299**:1221-1225.
126. Richter TA, Clisby KH: **Phenylthiocarbamide taste thresholds of rats and human beings.** *Am J Physiol* 1941, **134**:157-164.
127. Klein TW, DeFries JC: **Similar polymorphism of taste sensitivity to PTC in mice and men.** *Nature* 1970, **225**:555-557.
128. Whitney G, Harder DB: **PTC preference among laboratory mice: understanding of a previously "unreplicated" report.** *Behav Genet* 1986, **16**:605-610.
129. Smith JC, Davis JD, O-Keefe GB: **Lack of an order effect in brief contact taste tests with closely spaced test trials.** *Physiol Behav* 1992, **52**:1107-1111.
130. Spector AC: **Linking gustatory neurobiology to behavior in vertebrates.** *Neurosci Biobehav Rev* 2000, **24**:391-416.
131. Smith BK, Volaufova J, West DB: **Increased flavor preference and lick activity for sucrose and corn oil in SWR/J vs. AKR/J mice.** *Am J Physiol Regul Integr Comp Physiol* 2001, **281**:R596-606.
132. Boughter JD Jr, St John SJ, Noel DT, Ndubuizu O, Smith DV: **A brief-access test for bitter taste in mice.** *Chem Senses* 2002, **27**:133-142.
133. Nelson TM, Munger SD, Boughter JD Jr: **Taste sensitivities to PROP and PTC vary independently in mice.** *Chem Senses* 2003, **28**:695-704.
134. St John SJ, Pour L, Boughter JD Jr: **Phenylthiocarbamide produces conditioned taste aversions in mice.** *Chem Senses* 2005, **30**:377-382.
135. Whitney G, Harder DB: **Genetics of bitter perception in mice.** *Physiol Behav* 1994, **56**:1141-1147.
136. Lush IE, Hornigold N, King P, Stoye JP: **The genetics of tasting in mice. VII. Glycine revisited, and the chromosomal location of Sac and Soa.** *Genet Res* 1995, **66**:167-174.
137. Warren RP, Lewis RC: **Taste polymorphism in mice involving a bitter sugar derivative.** *Nature* 1970, **227**:77-78.
138. Lush IE: **The genetics of tasting in mice. I. Sucrose octaacetate.** *Genetical Research* 1981, **38**:93-95.
139. Lush IE: **The genetics of tasting in mice. III. Quinine.** *Genet Res* 1984, **44**:151-160.
140. Harder DB, Whitney G, Frye P, Smith JC, Rashotte ME: **Strain differences among mice in taste psychophysics of sucrose octaacetate.** *Chemical Senses* 1984, **9**:311-323.
141. Whitney G, Harder DB: **Single locus control of sucrose octaacetate tasting among mice.** *Behavior Genetics* 1986, **16**:559-574.
142. Harder DB, Capeless CG, Maggio JC, Boughter JD, Gannon KS, Whitney G, Azen EA: **Intermediate sucrose octa-acetate sensitivity suggests a third allele at mouse bitter taste locus Soa and Soa-Rua identity.** *Chemical Senses* 1992, **17**:391-401.
143. Inoue M, Li X, McCaughy SA, Beauchamp GK, Bachmanov AA: **Soa genotype selectively affects mouse gustatory neural responses to sucrose octaacetate.** *Physiological Genomics* 2001, **5**:181-186.
144. Capeless CG, Whitney G, Azen EA: **Chromosome mapping of Soa, a gene influencing gustatory sensitivity to sucrose octaacetate in mice.** *Behavior Genetics* 1992, **22**:655-663.
145. Bachmanov AA, Li X, Li S, Neira M, Beauchamp GK, Azen EA: **High-resolution genetic mapping of the sucrose octaacetate taste aversion (Soa) locus on mouse Chromosome 6.** *Mamm Genome* 2001, **12**:695-699.
146. Le Roy I, Pager J, Roubertoux PL: **Genetic dissection of gustatory sensitivity to bitterness (sucrose octaacetate) in mice.** *C R Acad Sci III* 1999, **322**:831-836.
147. Harder DB, Maggio JC, Whitney G: **Assessing gustatory detection capabilities using preference procedures.** *Chemical Senses* 1989, **14**:547-564.
148. Boughter JD Jr, Harder DB, Capeless CG, Whitney G: **Polygenic determination of quinine sensitivity among mice.** *Chem Senses* 1992, **17**:427-434.
149. Bachmanov AA, Reed DR, Tordoff MG, Price RA, Beauchamp GK: **Intake of ethanol, sodium chloride, sucrose, citric acid, and quinine hydrochloride solutions by mice: a genetic analysis.** *Behav Genet* 1996, **26**:563-573.
150. Harder DB, Whitney G: **A common polygenic basis for quinine and PROP avoidance in mice.** *Chem Senses* 1998, **23**:327-332.
151. Lush IE, Holland G: **The genetics of tasting in mice. V. Glycine and cycloheximide.** *Genet Res* 1988, **52**:207-212.
152. Blizard DA, Kotlus B, Frank ME: **Quantitative trait loci associated with short-term intake of sucrose, saccharin and quinine solutions in laboratory mice.** *Chem Senses* 1999, **24**:373-385.
153. Lush IE: **The genetics of tasting in mice. IV. The acetates of raffinose, galactose and β -lactose.** *Genetical Research* 1986, **47**:117-123.
154. Chandrashekar J, Mueller KL, Hoon MA, Adler E, Feng L, Guo W, Zuker CS, Ryba JP: **T2Rs function as bitter taste receptors.** *Cell* 2000, **100**:703-711.
155. Conte C, Guarin E, Marcuz A, Andres-Barquin PJ: **Functional expression of mammalian bitter taste receptors in *Caenorhabditis elegans*.** *Biochimie* 2006.
156. Go Y: **Lineage-specific expansions and contractions of the bitter taste receptor gene repertoire in vertebrates.** *Mol Biol Evol* 2006, **23**:964-972.
157. Conte C, Ebeling M, Marcuz A, Nef P, Andres-Barquin PJ: **Evolutionary relationships of the Tas2r receptor gene families in mouse and human.** *Physiol Genomics* 2003, **14**:73-82.
158. Wang X, Thomas SD, Zhang J: **Relaxation of selective constraint and loss of function in the evolution of human bitter taste receptor genes.** *Hum Mol Genet* 2004, **13**:2671-2678.
159. Go Y, Satta Y, Takenaka O, Takahata N: **Lineage-specific loss of function of bitter taste receptor genes in humans and nonhuman primates.** *Genetics* 2005, **170**:313-326.
160. Soranzo N, Bufe B, Sabeti PC, Wilson JF, Weale ME, Marguerie R, Meyerhof W, Goldstein DB: **Positive selection on a high-sensitivity allele of the human bitter-taste receptor TAS2R16.** *Curr Biol* 2005, **15**:1257-1265.
161. Gilbertson TA, Boughter JD Jr: **Taste transduction: appetizing times in gustation.** *NeuroReport* 2003, **14**:905-911.
162. Fitzsimons JT: *The Physiology of Thirst and Sodium Appetite* Cambridge: Cambridge University Press; 1979.
163. Denton D: *The hunger for salt: An anthropological, physiological and medical analysis* Berlin: Springer-Verlag; 1984.
164. Austin MA, King MC, Bawol RD, Hulley SB, Friedman GD: **Risk factors for coronary heart disease in adult female twins. Genetic heritability and shared environmental influences.** *Am J Epidemiol* 1987, **125**:308-318.
165. Hoshishima K, Yokoyama S, Seto K: **Taste sensitivity in various strains of mice.** *American Journal of Physiology* 1962, **202**:1200-1204.
166. Ninomiya Y, Sako N, Funakoshi M: **Strain differences in amiloride inhibition of NaCl responses in mice, *Mus musculus*.** *J Comp Physiol [A]* 1989, **166**:1-5.
167. Bachmanov AA, Schlager G, Tordoff MG, Beauchamp GK: **Consumption of electrolytes and quinine by mouse strains with different blood pressures.** *Physiol Behav* 1998, **64**:323-330.
168. Bachmanov AA, Beauchamp GK, Tordoff MG: **Voluntary consumption of NaCl, KCl, CaCl₂, and NH₄Cl solutions by 28 mouse strains.** *Behavior Genetics* 2002, **32**:445-457.
169. Beauchamp GK, Fisher AS: **Strain differences in consumption of saline solutions by mice.** *Physiol Behav* 1993, **54**:179-184.
170. Midkiff EE, Fitts DA, Simpson JB, Bernstein IL: **Absence of sodium chloride preference in Fischer-344 rats.** *American Journal of Physiology* 1985, **249**:R438-R442.

171. Midkiff EE, Bernstein IL, Fitts DA, Simpson JB: **Salt preference and salt appetite of Fischer-344 rats.** *Chem Senses* 1985, **10**:446.
172. Bachmanov AA: **The salt appetite of rats with spontaneous arterial hypertension.** *Fiziol Zh SSSR (in Russian)* 1989, **75**:942-947.
173. Tordoff MG: **Influence of dietary calcium on sodium and calcium intake of spontaneously hypertensive rats.** *American Journal of Physiology* 1992, **262**:R370-R381.
174. DiNicolantonio R, Mendelsohn FAO, Hutchinson JS: **Sodium chloride preference of genetically hypertensive and normotensive rats.** *American Journal of Physiology* 1983, **245**:R38-R44.
175. Sollars SI, Midkiff EE, Bernstein IL: **Genetic transmission of NaCl aversion in the Fisher-344 rat.** *Chemical Senses* 1990, **15**:521-527.
176. Bachmanov AA, Dmitriev YS: **Genetic analysis of some behavioral and physiological traits in hybrids between hypertensive and normotensive rats: Analysis of the inheritance nature (In Russian; English translation: Soviet Genetics, 28: 796-804).** *Genetika* 1992, **28**:150-159.
177. Bachmanov AA, Dmitriev YS: **Genetic analysis of some behavioral and physiological traits in hybrids between hypertensive and normotensive rats: Analysis of reciprocal differences (In Russian; English translation: Soviet Genetics, 28: 921-926).** *Genetika* 1992, **28**:120-127.
178. Bachmanov AA, Dmitriev YS: **Genetic analysis of some behavioral and physiological traits in hybrids between hypertensive and normotensive rats: Analysis of correlational associations (In Russian; English translation: Soviet Genetics, 28: 927-936).** *Genetika* 1992, **28**:128-138.
179. Harrap SB: **Genetic analysis of blood pressure and sodium balance in spontaneously hypertensive rats.** *Hypertension* 1986, **8**:572-582.
180. Di Nicolantonio R, Kren V, Zidek V, Pravenec M: **Salt preference of congenic strains derived from the spontaneously hypertensive rat.** *Physiol Behav* 2004, **80**:617-622.
181. Ninomiya Y, Bachmanov AA, Yatabe A, Beauchamp GK: **NaCl-preferring NZB/BINJ mice and NaCl-avoiding CBA/J mice have similar amiloride inhibition of chorda tympani responses to NaCl.** *Chemical Senses* 1998, **23**:411-415.
182. Ninomiya Y, Fukami Y, Yamazaki K, Beauchamp GK: **Amiloride inhibition of chorda tympani responses to NaCl and its temperature dependency in mice.** *Brain Research* 1996, **708**:153-158.
183. Gannon KS, Contreras RJ: **Sodium intake linked to amiloride-sensitive gustatory transduction in C57BL/6J and 129/J mice.** *Physiol Behav* 1995, **57**:231-239.
184. Formaker BK, Hill DL: **Lack of amiloride sensitivity in SHR and WKY glossopharyngeal taste responses to NaCl.** *Physiol Behav* 1991, **50**:765-769.
185. Minear MM, Hammack SE, Lundy RF Jr, Contreras RJ: **Amiloride inhibits taste nerve responses to NaCl and KCl in Sprague-Dawley and Fischer 344 rats.** *Physiol Behav* 1996, **60**:507-516.
186. Bernstein IL, Longley A, Taylor EM: **Amiloride sensitivity of chorda tympani response to NaCl in Fischer 344 and Wistar rats.** *Am J Physiol* 1991, **261**:R329-R333.
187. Brand J, Teeter J, Silver W: **Inhibition by amiloride of chorda tympani responses evoked by monovalent salts.** *Brain Research* 1985, **334**:207-214.
188. Eylam S, Spector AC: **Oral amiloride treatment decreases taste sensitivity to sodium salts in C57BL/6J and DBA/2J mice.** *Chem Senses* 2003, **28**:447-458.
189. Midkiff EE, Fitts DA, Simpson JB, Bernstein IL: **Attenuated sodium appetite in response to sodium deficiency in Fischer-344 rats.** *American Journal of Physiology* 1987, **252**:R562-R566.
190. Rowland NE, Fregly MJ: **Induction of an appetite for sodium in rats that show no spontaneous preference for sodium chloride solution - the Fischer 344 strain.** *Behavioral Neuroscience* 1988, **102**:961-968.
191. Caputo FA, Rowland NE, Fregly MJ: **Angiotensin-related intakes of water and NaCl in Fischer 344 and Sprague-Dawley rats.** *Am J Physiol* 1992, **262**:R382-388.
192. Grill HJ, Bernstein IL: **Strain differences in taste reactivity to NaCl.** *American Journal of Physiology* 1988, **255**:R424-R430.
193. Sollars SI, Sollars PJ, Bernstein IL: **Reversal of the sodium chloride aversion of Fischer 344 rats by chorda tympani nerve transection.** *Behav Neurosci* 1991, **105**:603-605.
194. Sollars SI, Bernstein IL: **Gustatory deafferentation and desalivation: effects on NaCl preference of Fischer 344 rats.** *Am J Physiol* 1994, **266**:R510-R517.
195. Clarke SN, Koh MT, Bernstein IL: **NaCl detection thresholds: comparison of Fischer 344 and Wistar rats.** *Chem Senses* 2001, **26**:253-257.
196. Formaker BK, Hill DL: **Peripheral taste responses in genetically hypertensive rats.** *Physiol Behav* 1990, **47**:1229-1237.
197. DiNicolantonio R, Hutchinson JS, Mendelsohn FA: **Exaggerated salt appetite of spontaneously hypertensive rats is decreased by central angiotensin-converting enzyme blockade.** *Nature* 1982, **298**:846-848.
198. Bachmanov AA, Whittlestone TH, Fitzsimons JT: **The enhanced sodium appetite of spontaneously hypertensive (SH) rats depends on angiotensin (Abstract).** *Journal of Physiology (London)* 1994, **480**:85P.
199. Drewnowski AJ, Brunzell JD, Sande K, Iverius PH, Greenwood MR: **Sweet tooth reconsidered: taste responsiveness in human obesity.** *Physiol Behav* 1985, **35**:617-622.
200. Salbe AD, DelParigi A, Pratley RE, Drewnowski A, Tataranni PA: **Taste preferences and body weight changes in an obesity-prone population.** *Am J Clin Nutr* 2004, **79**:372-378.
201. Yamamoto ME, Block GD, Ishii E: **Food patterns among adolescents: influence of alcohol consumption.** *Alcohol Clin Exp Res* 1991, **15**:359.
202. Kampov-Polevoy AB, Garbutt JC, Khalitov E: **Family history of alcoholism and response to sweets.** *Alcohol Clin Exp Res* 2003, **27**:1743-1749.
203. Kampov-Polevoy AB, Eick C, Boland G, Khalitov E, Crews FT: **Sweet liking, novelty seeking, and gender predict alcoholic status.** *Alcohol Clin Exp Res* 2004, **28**:1291-1298.
204. Kampov-Polevoy AB, Garbutt JC, Davis CE, Janowsky DS: **Preference for higher sugar concentrations and Tridimensional Personality Questionnaire scores in alcoholic and nonalcoholic men.** *Alcohol Clin Exp Res* 1998, **22**:610-614.
205. Kampov-Polevoy AB, Garbutt JC, Janowsky D: **Evidence of preference for a higher concentration sucrose solution in alcoholic men.** *American Journal of Psychiatry* 1997, **154**:269-270.
206. Guinard J-X, Zoumas-Morse C, Dietz J, Goldberg S, Holz M, Heck E, Amoros A: **Does consumption of beer, alcohol, and bitter substances affect bitterness perception.** *Physiol Behav* 1996, **59**:625-631.
207. Duffy VB, Davidson AC, Kidd JR, Kidd KK, Speed WC, Pakstis AJ, Reed DR, Snyder DJ, Bartoshuk LM: **Bitter receptor gene (TAS2R38), 6-n-propylthiouracil (PROP) bitterness and alcohol intake.** *Alcohol Clin Exp Res* 2004, **28**:1629-1637.
208. Intraruovo LR, Powers AS: **The perceived bitterness of beer and 6-n-propylthiouracil (PROP) taste sensitivity.** *Ann N Y Acad Sci* 1998, **855**:813-815.
209. Bartoshuk LM, Conner E, Grubin D, Karrer T, Kochenbach K, Palsco M, Snow D, Pelchat M, Danowski S: **PROP supertasters and the perception of ethyl alcohol.** *Chemical Senses* 1993, **18**:526-527.
210. Lanier SA, Hayes JE, Duffy VB: **Sweet and bitter tastes of alcoholic beverages mediate alcohol intake in of-age undergraduates.** *Physiol Behav* 2005, **83**:821-831.
211. Pelchat ML, Danowski S: **A possible genetic association between PROP-tasting and alcoholism.** *Physiol Behav* 1992, **51**:1261-1266.
212. DiCarlo ST, Powers AS: **Propylthiouracil tasting as a possible genetic association marker for two types of alcoholism.** *Physiol Behav* 1998, **64**:147-152.
213. Enoch MA, Harris CR, Goldman D: **Does a reduced sensitivity to bitter taste increase the risk of becoming nicotine addicted?** *Addict Behav* 2001, **26**:399-404.
214. Cannon DS, Baker TB, Piper ME, Scholand MB, Lawrence DL, Drayna DT, McMahon WM, Villegas GM, Caton TC, Coon H, Leppert MF: **Associations between phenylthiocarbamide gene polymorphisms and cigarette smoking.** *Nicotine Tob Res* 2005, **7**:853-858.
215. Dinehart ME, Hayes JE, Bartoshuk LM, Lanier SL, Duffy VB: **Bitter taste markers explain variability in vegetable sweetness, bitterness, and intake.** *Physiol Behav* 2006, **87**:304-313.
216. Basson MD, Bartoshuk LM, Dichello SZ, Panzini L, Weiffenbach JM, Duffy VB: **Association between 6-n-propylthiouracil (PROP) bitterness and colonic neoplasms.** *Dig Dis Sci* 2005, **50**:483-489.
217. Green BG: **The sensitivity of the tongue to ethanol.** *Annals of the New York Academy of Sciences* 1987, **510**:315-317.
218. Mattes RD, DiMeglio D: **Ethanol perception and ingestion.** *Physiol Behav* 2001, **72**:217-229.
219. Blizard DA: **Sweet and bitter taste of ethanol in C57BL/6J and DBA/2J mouse strains.** *Behav Genet* 2007, **37**:146-159.

220. Kiefer SW, Lawrence GJ: **The sweet-bitter taste of alcohol: aversion generalization to various sweet-quinine mixtures in the rat.** *Chemical Senses* 1988, **13**:633-641.
221. Kiefer SW, Mahadevan RS: **The taste of alcohol for rats as revealed by aversion generalization tests.** *Chemical Senses* 1993, **18**:509-522.
222. Molina JC, Domínguez HD, López MF, Pepino MY, Faas AE: **The role of fetal and infantile experience with alcohol in later recognition and acceptance patterns of the drug.** In *Alcohol and Alcoholism: Brain and Development* Edited by: Spear NE, Spear LP, Hannigan JH, Goodlett CR. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc; 1999:199-228.
223. Kiefer SW, Morrow NS: **Odor cue mediation of alcohol aversion learning in rats lacking gustatory neocortex.** *Behav Neurosci* 1991, **105**:25-32.
224. Nachman M, Larue C, Le Magnen J: **The role of olfactory and orosensory factors in the alcohol preference of inbred strains of mice.** *Physiol Behav* 1971, **6**:53-95.
225. Belknap JK, Belknap ND, Berg JH, Coleman R: **Preabsorptive vs. postabsorptive control of ethanol intake in C57BL/6J and DBA/2J mice.** *Behavior Genetics* 1977, **7**:413-425.
226. Blizard DA, McClearn GE: **Association between ethanol and sucrose intake in the laboratory mouse: exploration via congenic strains and conditioned taste aversion.** *Alcohol Clin Exp Res* 2000, **24**:253-258.
227. Lawrence GJ, Kiefer SW: **Generalization of specific taste aversions to alcohol in the rat.** *Chemical Senses* 1987, **12**:591-599.
228. Hellekant G, Danilova V, Roberts T, Ninomiya Y: **The taste of ethanol in a primate model: I. Chorda tympani nerve response in Macaca mulatta.** *Alcohol* 1997, **14**:473-484.
229. Sako N, Yamamoto T: **Electrophysiological and behavioral studies on taste effectiveness of alcohols in rats.** *American Journal of Physiology* 1999, **276**:R388-396.
230. Di Lorenzo PM, Kiefer SW, Rice AG, Garcia J: **Neural and behavioral responsivity to ethyl alcohol as a tastant.** *Alcohol* 1986, **3**:55-61.
231. Lemon CH, Brasser SM, Smith DV: **Alcohol activates a sucrose-responsive gustatory neural pathway.** *J Neurophysiol* 2004, **92**:536-544.
232. Levine AS, Kotz CM, Gosnell BA: **Sugars: hedonic aspects, neuroregulation, and energy balance.** *Am J Clin Nutr* 2003, **78**:834S-842S.
233. Gosnell BA, Majchrzak MJ: **Centrally administered opioid peptides stimulate saccharin intake in nondeprived rats.** *Pharmacol Biochem Behav* 1989, **33**:805-810.
234. George SR, Roldan L, Lui A, Naranjo CA: **Endogenous opioids are involved in the genetically determined high preference for ethanol consumption.** *Alcohol Clin Exp Res* 1991, **15**:668-672.
235. Hubell CL, Marglin SH, Spitalnic SJ, Abelson ML, Wild KD, Reid LD: **Opioidergic, serotonergic, and dopaminergic manipulations and rats' intake of a sweetened alcoholic beverage.** *Alcohol* 1991, **8**:355-367.
236. Pucilowski O, Rezvani AH, Janowsky DS: **Suppression of alcohol and saccharin preference in rats by a novel Ca²⁺ channel inhibitor, Goe 5438.** *Psychopharmacology* 1992, **107**:447-452.
237. Gentry RT, Dole VP: **Why does a sucrose choice reduce the consumption of alcohol in C57BL/6J mice?** *Life Sciences* 1987, **40**:2191-2194.
238. McMillen BA, Williams HL: **Role of taste and calories in the selection of ethanol by C57BL/6NHsd and Hsd:ICR mice.** *Alcohol* 1998, **15**:193-198.
239. Rodgers DA, McClearn GE, Bennett EL, Hebert M: **Alcohol preference as a function of its caloric utility in mice.** *J Comp Physiol Psychol* 1963, **56**:666-672.
240. Rodgers DA: **Factors underlying differences in alcohol preference among inbred strains of mice.** *Psychosomatic Medicine* 1966, **28**:498-513.
241. Freed EX: **Alcohol polydipsia in the rat as a function of caloric need.** *Quart J Stud Alc* 1972, **33**:504-507.
242. Richter CP: **Alcohol as a food.** *Quarterly Journal of Studies on Alcohol* 1941, **1**:650-662.
243. Overstreet DH, Kampov-Polevoy AB, Rezvani AH, Murelle L, Halikas JA, Janowsky DS: **Saccharin intake predicts ethanol intake in genetically heterogeneous rats as well as different rat strains.** *Alcohol Clin Exp Res* 1993, **17**:366-369.
244. Stewart RB, Russell RN, Lumeng L, Li TK, Murphy JM: **Consumption of sweet, salty, sour, and bitter solutions by selectively bred alcohol-preferring and alcohol-nonpreferring lines of rats.** *Alcohol Clin Exp Res* 1994, **18**:375-381.
245. Dess NK, Badia-Elder NE, Thiele TE, Kiefer SW, Blizard DA: **Ethanol consumption in rats selectively bred for differential saccharin intake.** *Alcohol* 1998, **16**:275-278.
246. Stewart RB, Bice P, Foroud T, Lumeng L, Li TK, Carr LG: **Correlation of saccharin and ethanol intake in the F2 progeny of HAD2 and LAD2 crosses.** *Alcohol Clin Exp Res* 2003, **27**:49A.
247. Blednov YA, Walker D, Martinez M, Levine M, Damak S, Margolskee RF: **Perception of sweet taste is important for voluntary alcohol consumption in mice.** *Genes Brain Behav* 2007.
248. Terenina-Rigaldie E, Moisan MP, Colas A, Beauge F, Shah KV, Jones BC, Mormede P: **Genetics of behaviour: phenotypic and molecular study of rats derived from high- and low-alcohol consuming lines.** *Pharmacogenetics* 2003, **13**:543-554.
249. Terenina-Rigaldie E, Jones BC, Mormede P: **Pleiotropic effect of a locus on chromosome 4 influencing alcohol drinking and emotional reactivity in rats.** *Genes Brain Behav* 2003, **2**:125-131.
250. Bufo B, Hofmann T, Krautwurst D, Raguse JD, Meyerhof W: **The human TAS2R16 receptor mediates bitter taste in response to beta-glucopyranosides.** *Nat Genet* 2002, **32**:397-401.
251. Kuhn C, Bufo B, Winnig M, Hofmann T, Frank O, Behrens M, Lewtschenko T, Slack JP, Ward CD, Meyerhof W: **Bitter taste receptors for saccharin and acesulfame K.** *J Neurosci* 2004, **24**:10260-10265.
252. Behrens M, Brockhoff A, Kuhn C, Bufo B, Winnig M, Meyerhof W: **The human taste receptor hTAS2R14 responds to a variety of different bitter compounds.** *Biochem Biophys Res Commun* 2004, **319**:479-485.
253. Bachmanov AA, Reed DR, Li X, Beauchamp GK: **Genetics of sweet taste preference.** *Pure and Applied Chemistry* 2002, **74**:1135-1140.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp