

Poster presentation

Open Access

## Transcriptional regulation network analysis of the hypertension-perturbed nucleus tractus solitarius

Gregory E Gonye\*, Rajanikanth Vadigepalli, Haiping Hao and James S Schwaber

Address: Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA

Email: Gregory E Gonye\* - [gonye@mail.dbi.tju.edu](mailto:gonye@mail.dbi.tju.edu)

\* Corresponding author

from Seventeenth Annual Computational Neuroscience Meeting: CNS\*2008  
Portland, OR, USA. 19–24 July 2008

Published: 11 July 2008

BMC Neuroscience 2008, 9(Suppl 1):P31 doi:10.1186/1471-2202-9-S1-P31

This abstract is available from: <http://www.biomedcentral.com/1471-2202/9/S1/P31>

© 2008 Gonye et al; licensee BioMed Central Ltd.

### Introduction

Neurons expressing the angiotensin II (angII) AT1 receptor (AT1R) in the nucleus tractus solitarius (NTS) have been implicated in many of the functions assigned to angII in the CNS including roles in neural control of cardiovascular function, in particular blood pressure regulation, the baroreflex, stress responses and maintenance of homeostasis (thirst and salt appetite). AngII, acting through the AT1 receptor has two main effects on neurons: a rapid and transient modulation of K<sup>+</sup> and Ca<sup>2+</sup> currents and longer lasting changes in gene expression, ultimately leading to modulation of noradrenergic neuronal networks regulating cardiovascular function [1]. Using a combined experimental and computational systems level analysis, we have investigated the transcriptional regulatory network downstream of AT1R activation by using N1E-115 neuroblastoma cells engineered to express AT1Rs as a model system and have recently extended this to the NTS.

### Methods

We have applied transcriptional regulatory network analysis (TRNA), using our bioinformatics framework, Promoter Analysis and Interaction Network Toolset (PAINT) [2] to the response of the NTS both directly to angII and to phenylephrine-induced acute hypertension. TRNA makes the fundamental assumption that many co-expressed genes share transcription regulatory elements (TRE's), typically transcription factor binding sites, in their promoters, leading to co-regulation. This results in a

higher frequency of the suspect binding sites in the promoters of sets of co-regulated genes relative to randomly associated sets of genes. PAINT uses bioinformatics in combination with robust statistical approaches to identify those regulatory elements significantly enriched in the promoters of the genes of interest relative to random gene sets. Predicted TF:promoter interactions are tested using chromatin immunoprecipitation PCR (ChIP-PCR) with chromatin from untreated and perturbed (angII or hypertension) NTS or cell lines.

### Results and conclusions

TRNA predictions generated from microarray-based gene expression analysis of stimulated NTS *ex vivo* by 100 nM angII for 4 hours (n = 4) are shown in Table 1. Predicted TF's include those previously described as angII-responsive in neurons (CREB, KROX), angII-responsive in non-neuronal cells (E2F, ATF and MEF2), and novel associations (Pax, cRel). Validation studies using chromatin immunoprecipitation (ChIP) to test specific TF::promoter interactions in the *in vivo* NTS and in an engineered AT1R-expressing neuronal cell line (N1E-AT1). Several predicted interactions have been confirmed, demonstrating both angII and hypertension-dependent binding dynamics. TRNA has successfully predicted TF activity dynamics at specific promoters relevant to neuronal function in hypertension and angII perturbed NTS. Going forward, these validated interactions will be manipulated *in vivo* to directly demonstrate a role in CNS cardiovascular regulation.

**Table 1: TRNA of ATIR-dependent NTS Expression**

TF Family	Enriched TRE	Fisher's p-value
CREB	V\$CREB_Q2_01	0.045
ATF	V\$ATF1_Q6	0.016
ATF	V\$ATF3_Q6	0.035
ATF	V\$ATF6_01	0.027
E2F	V\$E2F1_Q3_01	0.016
MEF2	V\$ZID_01	0.0065
Pax	V\$PAX3_B	0.0012
cRel	V\$CREL_01	0.0023
KROX/egr1	V\$KROX_Q6	0.036

### Acknowledgements

The authors wish to acknowledge the expert technical assistance of H. Liu, Y. Liu and M. Payne in support of these studies and the intellectual input of the other members of the Daniel Baugh Institute for Functional Genomics and Computational Biology. This work was supported by grants from the NIGMS and NHLBI (J. Schwaber, PI).

### References

1. Phillips MI, Summers C: **Angiotensin II in central nervous system physiology.** *Regul Pept* 1998, **78(1-3):1-11.**
2. Vadigepalli R, Chakravarthula P, Zak DE, Schwaber JS, Gonye GE: **PAINT: a promoter analysis and interaction network generation tool for gene regulatory network identification.** *Omics* 2003, **7(3):235-252.**

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

