

RESEARCH ARTICLE

Open Access



# Important modifications by sugammadex, a modified $\gamma$ -cyclodextrin, of ion currents in differentiated NSC-34 neuronal cells

Hung-Te Hsu<sup>1,2</sup>, Yi-Ching Lo<sup>3\*</sup>, Yan-Ming Huang<sup>4</sup>, Yu-Ting Tseng<sup>3,5</sup> and Sheng-Nan Wu<sup>4\*</sup>

## Abstract

**Background:** Sugammadex (SGX) is a modified  $\gamma$ -cyclodextrin used for reversal of steroidal neuromuscular blocking agents during general anesthesia. Despite its application in clinical use, whether SGX treatment exerts any effects on membrane ion currents in neurons remains largely unclear. In this study, effects of SGX treatment on ion currents, particularly on delayed-rectifier K<sup>+</sup> current [ $I_{K(DR)}$ ], were extensively investigated in differentiated NSC-34 neuronal cells.

**Results:** After cells were exposed to SGX (30  $\mu$ M), there was a reduction in the amplitude of  $I_{K(DR)}$  followed by an apparent slowing in current activation in response to membrane depolarization. The challenge of cells with SGX produced a depolarized shift by 15 mV in the activation curve of  $I_{K(DR)}$  accompanied by increased gating charge of this current. However, the inactivation curve of  $I_{K(DR)}$  remained unchanged following SGX treatment, as compared with that in untreated cells. According to a minimal reaction scheme, the lengthening of activation time constant of  $I_{K(DR)}$  caused by cell treatment with different SGX concentrations was quantitatively estimated with a dissociation constant of 17.5  $\mu$ M, a value that is clinically achievable. Accumulative slowing in  $I_{K(DR)}$  activation elicited by repetitive stimuli was enhanced in SGX-treated cells. SGX treatment did not alter the amplitude of voltage-gated Na<sup>+</sup> currents. In SGX-treated cells, dexamethasone (30  $\mu$ M), a synthetic glucocorticoid, produced little or no effect on L-type Ca<sup>2+</sup> currents, although it effectively suppressed the amplitude of this current in untreated cells.

**Conclusions:** The treatment of SGX may influence the amplitude and gating of  $I_{K(DR)}$  and its actions could potentially contribute to functional activities of motor neurons if similar results were found in vivo.

**Keywords:** Sugammadex, Motor neuron, Delayed-rectifier K<sup>+</sup> current, Activation kinetics, L-type Ca<sup>2+</sup> current, Glucocorticoid

## Background

Sugammadex (SGX) is recognized as a modified  $\gamma$ -cyclodextrin with a lipophilic core and a hydrophilic periphery, and it has been used clinically for reversal of neuromuscular blockade caused by rocuronium or vecuronium during general anesthesia [1–5]. A previous

report showed that addition of SGX could cause neuronal apoptosis in primary cultures [6]. This compound per se was also reported to be effective at reversing neurodegenerative disorder of the lower motor neurons [7, 8]. However, interestingly, how the treatment with SGX can perturb ionic currents remains largely unexplored.

The K<sub>v</sub>3 channels, a subfamily of K<sub>v</sub> channels, are distinguished from other types of K<sub>v</sub> channels by more positively shifted voltage-dependent activation and by faster activation and deactivation rates [9, 10]. These differences enable K<sub>v</sub>3 channels to be major determinants of high-frequency firing existing in several types of central

\*Correspondence: yichlo@kmu.edu.tw; snwu@mail.ncku.edu.tw

<sup>3</sup> Department of Pharmacology, School of Medicine, Kaohsiung Medical University, Kaohsiung City 80708, Taiwan

<sup>4</sup> Department of Physiology, National Cheng Kung University Medical College, No. 1, University Road, Tainan City 70101, Taiwan

Full list of author information is available at the end of the article



neurons [11–15]. The activity of  $K_v3.1$  channels has been recently described as playing a crucial role in controlling the amplitude of action potentials at synapses [16]. The de novo mutations in *KCNK1*, which encodes  $K_v3.1$  channels, have been also found to expand phenotypic spectra of this channel to progressive myoclonus epilepsy [17]. Therefore, these channels clearly are important targets used for investigations on electrical behaviors of central neurons including motor neurons [10].

The NSC-34 mouse motor neuron cell line is a hybridoma cell line derived from the fusion of mouse neuroblastoma with motor neuron-enriched spinal cord cells [18, 19]. These cells may create an easily accessible and clonally uniform motor neuron-like cell line that overcomes the problems associated with the culturing of primary spinal motor neurons. It has been demonstrated to be a suitable model for investigations on the mechanisms of neuronal development and differentiation in vitro and for studying electrophysiological properties of motor neurons in spite of being not considered as an adult motor neuron [18–20]. The biophysical properties of delayed-rectifier  $K^+$  current [ $I_{K(DR)}$ ] in NSC-34 cells were previously found to resemble the  $K_v3.1$ -encoded current because of positive mRNA detection of  $K_v3.1$  (*KCNK1*) [21, 22]. As NSC-34 cells were differentiated, the density of  $I_{K(DR)}$  was significantly enhanced. Previous work from our laboratory has shown that removal of membrane cholesterol by methyl- $\beta$ -cyclodextrin, a cyclic oligosaccharide, could modify activation kinetics of  $I_{K(DR)}$  in motor neuron-like cells [23]. Additionally, although voltage-gated  $Na^+$  current ( $I_{Na}$ ) in NSC-34 neuronal cells has been previously reported [24], few studies have been concerned with the biophysical or pharmacological properties of  $Ca^{2+}$  currents in these cells.

Therefore, the purpose of this work was to test whether SGX treatment could exert any perturbations on ionic currents present in NSC-34 neuronal cells differentiated with low serum and retinoic acid. The biophysical and pharmacological properties of ionic currents including delayed-rectifier  $K^+$  current [ $I_{K(DR)}$ ], voltage-gated  $Na^+$  current ( $I_{Na}$ ) and L-type  $Ca^{2+}$  current ( $I_{Ca,L}$ ) in untreated and SGX-treated cells have been characterized and compared in this study. Interestingly, the present results indicate that SGX treatment is capable of modifying the activation kinetics of  $I_{K(DR)}$  elicited by membrane depolarization in these cells in a concentration-, time-, and state-dependent manner.

## Methods

### Drugs and solutions

Sugammadex (Bridion<sup>®</sup>, SGX,  $C_{72}H_{104}Na_8O_{48}S_8$ ) was obtained from Schering-Plough (Kenilworth, NJ, USA), aconitine, dexamethasone (DEX), L-aspartic acid,

neostigmine, nifedipine, retinoic acid, tetraethylammonium chloride (TEA) and tetrodotoxin were from Sigma-Aldrich (St. Louis, MO, USA), iberiotoxin, apamin and  $\omega$ -conotoxin GVIA were from Alomone Labs (Jerusalem, Israel), isobavachalcone (2',4',4'-trihydroxy-3'-[3''-methylbut-3''-enyl]chalcone) was from Enzo Life Sci. (Plymouth Meeting, PA, USA), midazolam was from Nang Kuang Pharmaceutical Co. (Tainan City, Taiwan) and ranolazine [(±)-*N*-(2,6-dimethyl-phenyl)-4[2-hydroxy-3(2-methoxy-phenoxy)propyl]-1-piperazine acetamide] was from Tocris Cookson, Ltd. (Bristol, UK). For cell preparations, all culture media, fetal bovine serum (FBS), L-glutamine, trypsin/EDTA, penicillin–streptomycin and amphotericin B were obtained from Invitrogen (Carlsbad, CA, USA). All other chemicals including  $CdCl_2$ , CsCl, CsOH and  $MgSO_4$  were obtained from regular commercial chemicals and of reagent grade. The compositions of bathing and pipette solutions used in this study are illustrated in Table 1.

### Cell preparation and differentiation

NSC-34 neuronal cells were originally produced by fusion of the motor neuron-enriched, embryonic mouse spinal cords with the mouse neuroblastoma [19]. These cells were kindly provided by Professor Yuh-Jyh Jong (Department of Pediatrics, Kaohsiung Medical University Hospital, Kaohsiung City, Taiwan). They were routinely grown in 1:1 mixture of Dulbecco's modified Eagle medium (DMEM) and Ham's F12 medium that was supplemented with 10% (v/v) FBS and 1% penicillin–streptomycin. Cultures were incubated at 37 °C in a humidified environment of 5%  $CO_2$ /95% air. The medium was replenished every 2–3 days for removal of non-adhering cells. To slow cell proliferation and enhance their maturation towards a differentiated state [20], before confluence, cells were grown in 1:1 DMEM plus Ham's F12 medium supplemented with low serum (1% FBS) and 1  $\mu$ M retinoic acid. The SGX-treated cells were incubated at 37 °C for 1 h in normal Tyrode's solution containing different concentrations of this compound. The reason that the duration was set at 1 h was to ensure that SGX could exert its interaction with cell membrane.

### Electrophysiological measurements

Shortly before each experiment, cells were dissociated, and an aliquot of cell suspension was transferred to a homemade recording chamber positioned on the stage of a CKX-41 inverted microscope (Olympus, Tokyo, Japan). Cells were immersed at room temperature (20–25 °C) in normal Tyrode's solution containing 1.8 mM  $CaCl_2$ . The patch electrodes used were prepared from Kimax capillary tubes (#34500; Kimble Glass, Vineland, NJ, USA) using a vertical two-step electrode puller

**Table 1 Composition of normal Tyrode's solution and the pipette solution used in this study**

Solution	Purpose or name	Composition
Bathing solution	Normal Tyrode's solution	136.5 mM NaCl, 5.4 mM KCl, 1.8 mM CaCl <sub>2</sub> , 0.53 mM MgCl <sub>2</sub> , 5.5 mM glucose, and 5.5 mM HEPES–NaOH buffer, pH 7.4
Pipette solution	For recordings of K <sup>+</sup> current or membrane potential	130 mM K-aspartate, 20 mM KCl, 1 mM KH <sub>2</sub> PO <sub>4</sub> , 1 mM MgCl <sub>2</sub> , 3 mM Na <sub>2</sub> ATP, 0.1 mM Na <sub>2</sub> GTP, 0.1 mM EGTA, and 5 mM HEPES–KOH buffer, pH 7.2
	For recordings of Na <sup>+</sup> or Ca <sup>2+</sup> currents	130 mM Cs-aspartate, 20 mM CsCl, 1 mM KH <sub>2</sub> PO <sub>4</sub> , 1 mM MgCl <sub>2</sub> , 3 mM Na <sub>2</sub> ATP, 0.1 mM Na <sub>2</sub> GTP, 0.1 mM EGTA, and 5 mM HEPES–CsOH buffer, pH 7.2

(PP-83 or PP-830; Narishige, Tokyo, Japan), and their tips were then fire-polished with an MF-83 micro-forge (Narishige). Experiments were performed using the whole-cell configuration of standard patch-clamp technique using either an RK-400 (Bio-Logic, Claix, France) or an Axopatch 200B (Molecular Devices, Sunnyvale, CA, USA) patch-clamp amplifier [19]. Junctional potentials that developed when the composition of the pipette solution was different from that in the bath were nulled.

#### Data recordings

The signals consisting of voltage and current tracings were displayed and recorded online using an ASUSPRO-BU401LG computer (ASUS, Taipei City, Taiwan) equipped with a Digidata 1440A device (Molecular Devices), and the experiments were controlled by pCLAMP 10.2 software (Molecular Devices). Current signals were low-pass filtered at 3 kHz and digitized at 10 kHz. In some experiments of verifying analog-to-digital conversion, signals are digitized using a PowerLab acquisition system with LabChart 7.0 programs (AD Instruments, Gerin, Tainan City, Taiwan). The resultant data achieved during this experiment were analyzed off-line by use of various analytical tools including the LabChart 7.0 program (Gerin), Origin 8.0 (OriginLab, Northampton, MA, USA) and custom-made macro procedures run under Excel 2013 (Microsoft, Redmond, WA, USA). The voltage-step profiles digitally created from pCLAMP 10.2 were employed to evaluate current–voltage ( $I$ – $V$ ) relationships or steady-state inactivation of ionic currents [e.g.,  $I_{K(DR)}$ ].

#### Data analyses

The relationships between the relative  $I_{K(DR)}$  amplitude and the membrane potential obtained with or without the treatment of SGX (30  $\mu$ M) were fitted with a Boltzmann function of the following form:

$$\frac{I}{I_{\max}} = \frac{1}{1 + \exp\left[\frac{-(V - V_{1/2})qF}{RT}\right]},$$

where  $I_{\max}$  is the maximal amplitude of  $I_{K(DR)}$  elicited by membrane depolarization,  $V_{1/2}$  the voltage at which there

is half-maximal activation,  $q$  the apparent gating charge,  $F$  Faraday's constant,  $R$  the universal gas constant, and  $T$  the absolute temperature.

The free energy involved in the gating of  $I_{K(DR)}$  ( $\Delta G_0$ ) was calculated assuming that there is a 2-state (i.e., closed [resting] and open) gating model inherently in the channel.  $\Delta G_0$  for  $I_{K(DR)}$  activation at 0 mV with or without treatment of SGX would be equal to  $q \times F \times V_{1/2}$  [25, 26]. The standard errors of  $\Delta G_0$  (i.e.,  $\sigma_{qFV_{1/2}}$ ) were calculated according to:

$$\sigma_{qFV_{1/2}} = F \times \sqrt{V_{1/2}^2 \sigma_q^2 + q^2 \sigma_{V_{1/2}}^2}$$

where  $\sigma_q$  and  $\sigma_{V_{1/2}}$  represent the standard errors in  $q$  and  $V_{1/2}$  respectively.

Perturbation by SGX treatment of free energy involved in the gating of  $I_{K(DR)}$  was calculated as  $\Delta \Delta G_0 = \Delta G_0^{SGX} - \Delta G_0^{Ctrl} = F(qV_{1/2SGX} - qV_{1/2Ctrl}) = \Delta(qFV_{1/2})$ , where  $\Delta G_0^{Ctrl}$  and  $\Delta G_0^{SGX}$  indicate the free energy of  $I_{K(DR)}$  activation taken from untreated cells and cells exposed to SGX respectively.

The steady-state inactivation curve of  $I_{K(DR)}$  with or without treatment of SGX was derived and plotted against the conditioning pulses and then fitted to another Boltzmann equation:

$$\frac{I}{I_{\max}} = \frac{1}{1 + \exp\left[\frac{(V - V_{1/2})}{k}\right]},$$

where  $V$  represents the conditioning potential in mV,  $V_{1/2}$  is the membrane potential for half-maximal inactivation, and  $k$  is the slope factor of inactivation curve for  $I_{K(DR)}$  elicited by membrane depolarization.

Linear or nonlinear curve-fitting to data sets presented herein was performed using either Microsoft Solver function embedded in Excel (Microsoft) or Origin 8.0 program (OriginLab). The values are provided as the mean  $\pm$  standard error of the mean (SEM) with sample sizes ( $n$ ) indicating the cell number from which the results were obtained. The paired or unpaired Student's  $t$  test and a one-way analysis of variance with the least-significant difference method for multiple comparisons

were used for statistical evaluation of differences among means. Statistical analyses were performed using the Statistical Package for Social Science 20 (SPSS; IBM Corp., Armonk, New York). Statistical significance was determined at a  $P$  value of  $<0.05$ .

## Results

### Effect of SGX treatment on delayed-rectifier $K^+$ currents [ $I_{K(DR)}$ ] in differentiated NSC-34 neuronal cells

In the first set of experiments, the whole-cell configuration of the patch-clamp technique was conducted to investigate effects of SGX on ionic currents in these cells. Cells were bathed in  $Ca^{2+}$ -free Tyrode's solution that contained tetrodotoxin (1  $\mu M$ ) and 0.5 mM  $CdCl_2$ , and the recording pipette was filled with  $K^+$ -containing solution described in "Methods". Figure 1A depicts the  $I_{K(DR)}$  amplitudes when cells were exposed to different concentrations (10, 30 and 100  $\mu M$ ) of SGX. Because the SGX concentration at 30  $\mu M$  was close to the  $IC_{50}$  value for inhibition of  $I_{K(DR)}$ , most experiments shown below used such concentrations. As illustrated in Fig. 1B, C, as the cell was held at  $-50$  mV and different voltage steps ranging from  $-50$  to  $+60$  mV in 10-mV increments were applied, a family of large outward currents was readily elicited. Current amplitudes were increased with greater depolarizations in an outward-rectifying manner with a reversal potential of  $-76 \pm 2$  mV ( $n = 12$ ). These outward currents, which were insensitive to inhibition by either iberiotoxin (200 nM) or apamin (200 nM), have been referred to as  $I_{K(DR)}$ , the biophysical properties of which resemble the  $K_{V3.1}$ -encoded  $K^+$  currents [12, 22]. In particular, as the cells were exposed to SGX (30  $\mu M$ ), the amplitude and gating of  $I_{K(DR)}$  were altered throughout the entire range of voltage-clamp steps as compared with those taken from untreated cells. For example, when the voltage step from  $-50$  to  $+50$  mV was evoked, SGX (10  $\mu M$ ) treatment significantly decreased the amplitude of initial  $I_{K(DR)}$  (i.e., the 60th millisecond after the beginning of voltage step) by  $41 \pm 3\%$  from  $1059 \pm 207$  to  $625 \pm 167$  pA ( $n = 11$ ,  $P < 0.05$ ). However, such treatment slightly but significantly decreased  $I_{K(DR)}$  amplitude at the end of depolarizing pulse by  $22 \pm 2\%$  from  $976 \pm 197$  to  $766 \pm 89$  pA ( $n = 11$ ,  $P < 0.05$ ). No significant difference in cell capacitance between untreated ( $17.9 \pm 1.8$  pF,  $n = 13$ ) and SGX-treated ( $18.1 \pm 1.9$  pF,  $n = 13$ ,  $P > 0.05$ ) cells could be clearly demonstrated. The averaged amplitude versus voltage relationships of  $I_{K(DR)}$  were measured and plotted in untreated and SGX-treated cells (Fig. 1C). The  $I-V$  relationships with or without SGX treatment were then obtained at the beginning [Fig. 1C(a)] and end [Fig. 1C(b)] of voltage pulses; therefore, SGX suppressed  $I_{K(DR)}$  amplitude in a concentration-dependent manner. Additionally, neostigmine (1  $\mu M$ ), a typical drug used for

reversing the effect of rocuronium-induced neuromuscular blockade, had no effect on the amplitude of gating of  $I_{K(DR)}$  in these cells (Additional file 1: Fig. S1).


### The activation curve of $I_{K(DR)}$ obtained with or without treatment of SGX

Figure 2 shows the activation curve of  $I_{K(DR)}$  in the absence and presence of SGX (30  $\mu M$ ) treatment. The plot of relative  $I_{K(DR)}$  amplitude as a function of membrane potential was determined and fitted with a Boltzmann function as described under "Methods". In untreated cells,  $V_{1/2} = 21.1 \pm 2.2$  mV and  $q = 1.93 \pm 0.04 e$  ( $n = 9$ ), while in SGX-treated cells,  $V_{1/2} = 36.1 \pm 1.9$  mV and  $q = 2.98 \pm 0.17 e$  ( $n = 9$ ). The data showed that, as differentiated NSC-34 cells were treated with SGX (30  $\mu M$ ), the activation curve of this current was shifted along the voltage axis to more positive potentials by approximately 15 mV and the elementary charge for activation was elevated 1.5-fold.


According to the values of  $q$  and  $V_{1/2}$ , the free energy involved in the gating of  $I_{K(DR)}$  at 0 mV ( $\Delta G_0$ ) in the absence and presence of SGX treatment was estimated to be  $3.93 \pm 0.12$  and  $10.35 \pm 0.18$  kJ/mol ( $n = 9$ ) respectively. The perturbation by SGX treatment of free energy ( $\Delta\Delta G_0$ ) involved in  $I_{K(DR)}$  gating was calculated to be  $6.42 \pm 0.15$  kJ/mol. It is therefore anticipated from these data that SGX treatment can increase the free energy needed for  $I_{K(DR)}$  activation observed in differentiated NSC-34 cells.

### Kinetic evaluation of $I_{K(DR)}$ block by SGX

During cell exposure to SGX, the activation time course of  $I_{K(DR)}$  in response to membrane depolarization tended to become slower. The activation kinetics of SGX-induced block of  $I_{K(DR)}$  in response to membrane depolarization was further quantitatively evaluated in cells exposed to different SGX concentrations. The concentration dependence of  $I_{K(DR)}$  inhibition by SGX treatment is illustrated in Fig. 3. The results showed that its effects on  $I_{K(DR)}$  could exert a concentration-dependent increase in the rate of development of inhibition. Consequently, this effect on  $I_{K(DR)}$  in differentiated NSC-34 cells can be explained by a state-dependent blocking mechanism in which this compound may preferentially bind to the closed (resting) state of the  $K_V$  channels according to a minimal kinetic scheme:



where  $\alpha$  and  $\beta$  represent kinetic constants for the opening and closing of the  $K_V$  channel respectively;  $k_{+1}$  and  $k_{-1}$  are those for blocking and unblocking by SGX treatment; and  $[B]$  is the SGX concentration.  $C$ ,  $O$  and  $C \cdot B$




**Fig. 1** Effect of SGX treatment on  $I_{K(DR)}$  in differentiated NSC-34 neuronal cells. In these experiments, cells were bathed in  $Ca^{2+}$ -free Tyrode's solution containing 1  $\mu$ M tetrodotoxin and the recording pipette was filled with  $K^+$ -containing solution. The treated cells were incubated with SGX (30  $\mu$ M) for 1 h at 37  $^{\circ}C$ . **(A)** Bar graph showing the data of  $I_{K(DR)}$  amplitude when cells were treated with 10, 30 and 100  $\mu$ M SGX. Current amplitudes were measured at the beginning of depolarizing voltage was obtained at the 60th milliseconds after the initial rise of voltage from  $-50$  to  $+50$  mV (mean  $\pm$  SEM;  $n = 10-12$  for each bar). \*Significantly different from control ( $P < 0.05$ ). **(B)** Superimposed current traces obtained in untreated (upper) and SGX-treated (lower) cells. The cells examined were held at  $-50$  mV and the voltages ranging from  $-50$  to  $+60$  mV in 10-mV increments were applied, as whole-cell recordings were established. The uppermost part indicates the voltage protocol used. **(C)** Current amplitude versus membrane potential relationships of  $I_{K(DR)}$  in untreated cells (square symbols) and in cells treated with 30  $\mu$ M SGX (circle symbols). In **C(a)** and **C(b)**,  $I_{K(DR)}$  amplitude was measured at the beginning (filled symbols) and end (open symbols) of depolarizing steps, respectively.  $I_{K(DR)}$  amplitudes measured at the beginning of depolarizing voltage were obtained at the 60th milliseconds after the initial rise of voltage. Each point represents the mean  $\pm$  SEM ( $n = 10-12$ ). \*Significantly different from controls [i.e.,  $I_{K(DR)}$  amplitude at the same level of voltage step] ( $P < 0.05$ )

indicate the closed (resting), open, and closed-blocked state respectively.

Blocking and unblocking rate constants,  $k_{+1}$  and  $k_{-1}$ , could be determined from activation time constants of

SGX-sensitive  $I_{K(DR)}$  (i.e., difference in  $I_{K(DR)}$  taken from untreated and SGX-treated cells) obtained in different concentrations of this compound. The rate constants were then computed using the relation:





**Fig. 2** Effect of SGX on the activation curve of  $I_{K(DR)}$  recorded from differentiated NSC-34 neuronal cells. In these experiments, cells were bathed in  $Ca^{2+}$ -free Tyrode's solution. Voltage dependence of  $I_{K(DR)}$  activation in the absence (filled square symbol) and presence (open square symbol) of SGX (30  $\mu$ M) treatment (mean  $\pm$  SEM;  $n = 9$  for each point) is illustrated. The smooth lines obtained in untreated and SGX-treated cells represent the best fits to the Boltzmann equation described in "Methods". Notably, SGX treatment can shift the activation curve of  $I_{K(DR)}$  to a depolarizing voltage, together with increase in the gating charge of this current

$$\frac{1}{\tau_{act}} = k_{+1}[B] + k_{-1},$$

where  $k_{+1}$  and  $k_{-1}$  respectively resulted from the slope and from the y-axis intercept at  $[B] = 0$  of the linear regression interpolating the reciprocal time constants ( $1/\tau_{act}$ ) versus the SGX concentration. The relationship between  $1/\tau_{act}$  and  $[B]$  was computed to be linear with a correlation coefficient of 0.95 (Fig. 3B). The blocking and unblocking rate constants were thereafter calculated to be  $0.0012 \text{ ms}^{-1} \mu\text{M}^{-1}$  and  $0.021 \text{ ms}^{-1}$  respectively. According to these rate constants, dividing  $k_{-1}$  by  $k_{+1}$  yielded a dissociation constant ( $K_D$ ) of  $17.5 \mu\text{M}$  under SGX treatment.

#### Inability of SGX to modify the steady-state inactivation curve of $I_{K(DR)}$

In order to further characterize effect of SGX treatment on  $I_{K(DR)}$ , we next studied the inactivation of  $I_{K(DR)}$  in differentiated cells using a two-step voltage protocol. The quasi-inactivation curves of  $I_{K(DR)}$  in untreated and SGX-treated cells are illustrated in Fig. 4. In this set of experiments, a 10-s conditioning pulse in different membrane potentials preceded a test potential (300 ms in duration) to +80 mV from a holding potential of -90 mV (Fig. 4a). The relationships between the conditioning pulses and the normalized amplitudes of  $I_{K(DR)}$  in untreated and SGX-treated cells were constructed and fitted by the Boltzmann equation described in "Methods". In untreated cells,  $V_{1/2} = 6.2 \pm 0.9 \text{ mV}$  and  $k = 0.42 \pm 0.02 \text{ mV}$  ( $n = 9$ ), while in SGX-treated cells,


$V_{1/2} = 6.3 \pm 0.09 \text{ mV}$  and  $k = 0.44 \pm 0.03 \text{ mV}$  ( $n = 9$ ). Distinguishable from the results of  $I_{K(DR)}$  activation curve obtained in untreated and SGX-treated cells, the values for both  $V_{1/2}$  and  $k$  were not found to differ significantly between the two groups of cells ( $P > 0.05$ ). The experimental results therefore suggest that there is little or no modification in the voltage-dependent profile of  $I_{K(DR)}$  inactivation following SGX treatment.

#### The increase of cumulative inhibition of $I_{K(DR)}$ activation in SGX-treated cells

In another set of experiments, we sought to determine whether in SGX-treated cells, the activation time course of  $I_{K(DR)}$  induced by repetitive depolarizing stimuli could be altered. Under control conditions (i.e., in untreated cells), a single 100-ms depolarizing step to +50 mV from a holding potential of -50 mV was applied to produce an exponential increase of  $I_{K(DR)}$  with a time constant of  $27 \pm 2 \text{ ms}$  ( $n = 10$ ). However, the activation time constant for 10-ms repetitive pulses to +50 mV, each of which lasted 10 ms with 5-ms interval at -50 mV between the depolarizing pulses, was significantly reduced to  $19 \pm 2 \text{ ms}$  ( $n = 10$ ,  $P < 0.05$ ). The results indicate a progressive increase in the activation rate of  $I_{K(DR)}$  in response to repetitive depolarizing stimuli (Fig. 5 and Additional file 2: Fig. S2). However, in SGX-treated cells, the value of activation time constant obtained during this train of short repetitive pulses became raised. In the cells exposed to 30  $\mu\text{M}$  SGX, the time constants were significantly increased to  $47 \pm 5 \text{ ms}$  ( $n = 9$ ,  $P < 0.05$ ); accordingly, the results showed that there was an excessive accumulative slowing in the activation of  $I_{K(DR)}$  as cells were exposed to SGX.

#### Effect of SGX treatment on voltage-gated $Na^+$ current ( $I_{Na}$ ) in differentiated NSC-34 cells


In the next set of experiments, we investigated effect of SGX treatment on  $I_{Na}$  in these cells. Untreated or SGX-treated cells were bathed in  $Ca^{2+}$ -free Tyrode's solution containing 10 mM TEA and the recording pipette was filled with  $Cs^+$ -containing solution, the composition of which is shown in Table 1. The  $I_{Na}$  was elicited by depolarizing pulse from -80 to -10 mV with a duration of 100 ms. In our experiments, SGX treatment at 10, 30, and 100  $\mu\text{M}$  did not exert any significant effect on the peak amplitude of  $I_{Na}$  or  $I_{Ca,L}$  in these cells. As shown in Fig. 6, there was no significant difference in the amplitude of  $I_{Na}$  between untreated cells ( $1.82 \pm 0.3 \text{ nA}$ ,  $n = 8$ ) and cells treated with SGX (10  $\mu\text{M}$ ) ( $1.81 \pm 0.2 \text{ nA}$ ,  $n = 8$ ,  $P > 0.05$ ). However, in SGX-treated cells, addition of ranolazine (10  $\mu\text{M}$ ), a blocker of  $I_{Na}$  [27], was capable of suppressing the peak amplitude of  $I_{Na}$  significantly from





$1.82 \pm 0.2$  nA to  $0.98 \pm 0.1$  nA ( $n = 7$ ,  $P < 0.05$ ). Therefore, SGX treatment per se did not modify the amplitude of  $I_{Na}$  in these cells, while the presence of ranolazine remained effective at suppressing  $I_{Na}$  in SGX-treated cells.

#### Comparison of the effect of dexamethasone (DEX) on L-type $\text{Ca}^{2+}$ current ( $I_{Ca,L}$ ) in SGX-treated cells

SGX is a modified  $\gamma$ -cyclodextrin and able to encapsulate steroid-like compounds [1, 4]. DEX, a synthetic glucocorticoid, can produce an inhibitory effect on the reversal of rocuronium-induced neuromuscular block by SGX in functionally innervated human muscle cells [28]. Therefore, in a final set of experiments, it was further examined whether the amplitude of  $I_{Ca,L}$  recorded from SGX-treated cells could be altered by DEX. The experiments were performed as cells were bathed in normal Tyrode's



solution containing  $1.8 \text{ mM}$   $\text{CaCl}_2$  and  $1 \mu\text{M}$  tetrodotoxin, and the recording pipette was filled with  $\text{Cs}^+$ -containing solution. The peak amplitude of  $I_{Ca,L}$  in response to membrane depolarization from  $-50$  to  $0$  mV was effectively suppressed by the presence of nifedipine ( $1 \mu\text{M}$ ), a blocker of  $I_{Ca,L}$ , but not by  $\omega$ -conotoxin GVIA ( $1 \mu\text{M}$ ), a toxin isolated from *Conus geographus* and known to block N-type  $\text{Ca}^{2+}$  current (Additional file 3: Fig. S3). As illustrated in Fig. 7, as untreated cells were depolarized from  $-50$  to  $0$  mV, DEX ( $30 \mu\text{M}$ ) significantly suppressed the peak amplitude of  $I_{Ca,L}$  from  $51.7 \pm 5.3$  to  $24.5 \pm 2.6$  pA ( $n = 11$ ,  $P < 0.05$ ); however, the overall  $I-V$  relationship of this current remained unchanged in the presence of DEX. The concentration of DEX ( $30 \mu\text{M}$ ) used in this study was fundamentally based on a previous report [29]. The results are compatible with previous observations made in pituitary tumor cells [29]. In contrast, in SGX-treated cells, DEX at the same concentration had no significant effect on the amplitude of  $I_{Ca,L}$  [ $52.1 \pm 5.3$  pA (control),  $n = 11$  versus  $51.9 \pm 5.4$  pA (in the presence of DEX),  $n = 11$ ,  $P > 0.05$ ]; therefore, findings from these



results indicate that, in SGX-treated cells, DEX-mediated inhibition of  $I_{Ca,L}$  in response to membrane depolarization was abolished.


### Discussion

In this study, the blocking of  $I_{K(DR)}$  by SGX treatment was noted to be not instantaneous, but to develop with time immediately after the cell became depolarized. Such treatment produces a time-dependent increase in the activation time constant of  $I_{K(DR)}$  in response to membrane depolarization. However, the inactivation curve of  $I_{K(DR)}$  obtained in untreated and SGX-treated cells did not differ significantly. It thus appears that, as cells are exposed to SGX, this compound has a greater affinity for the closed (resting) state in the  $K_V$  channel existing on differentiated NSC-34 cells. The activated channels during SGX treatment tended to produce a lower affinity than those residing in the closed state. As a result, the transition from closed to open state became slowed during cell exposure to SGX. Based on our study, it is therefore tempting to speculate that the treatment of SGX or other structurally similar agents binds to the closed state of the channel and/or blocks a prolonged channel closing in  $K_V3.1$  channels. Alternatively, the challenge of cells with SGX or other structurally similar agents can modulate  $K_V3.1$  channels where the closed channel conformation represents the high-affinity binding site. It also needs to be noted that, according to minimal reaction scheme shown here, the dissociation constant for SGX treatment was calculated to be  $17.5 \mu M$ , a value that can be clinically achievable [1, 3, 5].

The  $K_V3.1$ -encoded currents were reported to be the major molecular components of  $I_{K(DR)}$  in these cells [20, 21]. Importantly, our study demonstrated that the activation kinetics of  $I_{K(DR)}$  (i.e.,  $K_V3.1$ -encoded current) in SGX-treated cells virtually became slowed in a time- and state-dependent manner. The results were in contrast with inhibitory effects of midazolam or aconitine on  $I_{K(DR)}$  [12, 21]. These two compounds exerted inhibitory effects via a mechanism through binding to the open state of the channel followed by increased rate of  $I_{K(DR)}$  inactivation. During repetitive stimuli, the perturbation by SGX treatment of  $I_{K(DR)}$  activation was potentiated.

By virtue of computational analysis, previous work has shown that changes in activation kinetics of  $I_{K(DR)}$  might lead to generation of action potentials with spike-frequency adaptation [23]. The ability of  $K_V3.1$ -encoded currents to control the waveforms of action potentials at synapses has recently been reported [16]. Indeed, different de novo mutations in *KCNKI* have been reported to display a wide variety of progressive myoclonus epilepsy [17]. However, SGX treatment had little or no effect on





the peak amplitude of  $I_{\text{Na}^+}$ . Therefore, the present results showing any changes in the amplitude and gating by SGX treatment of  $I_{\text{K(DR)}}$  can be of pharmacological and clinical relevance.

Following SGX treatment,  $I_{\text{K(DR)}}$  enriched in differentiated NSC-34 cells became activated at more depolarized voltages in comparison with that from untreated cells. Moreover, the steepness of activation curve for  $I_{\text{K(DR)}}$  became significantly greater in cells exposed to SGX, indicating that the effective number of elementary charges during channel activation in SGX-treated cells was significantly raised. These results are important because they led us to estimate that energy change ( $\Delta G_0^{\text{SGX}}$ ) for generation of  $I_{\text{K(DR)}}$  was a value of 10.35 kJ/mol. This value was found to be significantly greater than that  $\Delta G_0^{\text{Ctrl}}$  (i.e., 3.93 kJ/mol) in untreated cells. SGX treatment apparently is involved in voltage-sensitive gating functions of  $I_{\text{K(DR)}}$ , despite no clear change in inactivation curve of  $I_{\text{K(DR)}}$  between the two groups of cells. The results lead us to propose that following SGX treatment, the energy barrier for activation of  $\text{K}_v3.1$  channels became elevated.

In our experimental conditions, supplementation of the medium with retinoic acid resulted in changes in cell

morphology and an increase in mRNA expression of the  $\text{K}_v3.1$  subunit in differentiated NSC-34 neuronal cells [20, 21]. However, the modification of  $I_{\text{K(DR)}}$  kinetics by SGX presented here did not appear to occur by the gene regulation of these channels, because significant changes in this current in differentiated NSC-34 cells generally occurred with a short time course. Moreover, no changes in  $I_{\text{K(DR)}}$  density after treatment with SGX were observed, suggesting that such maneuver did not alter the main parts of ion channel permeation pathway (i.e., the S5 and S6 regions). It is thus possible that SGX treatment can regulate the gating kinetics of  $I_{\text{K(DR)}}$  with no apparent change in the number of functional channels on plasma membrane.

Consistent with previous studies [29], we clearly demonstrated that addition of DEX suppressed the peak amplitude of  $I_{\text{Ca,L}}$  in differentiated NSC-34 neuronal cells. It is important to note, however, that the inhibition by DEX of  $I_{\text{Ca,L}}$  did not occur in SGX-treated cells, despite the ability of  $\text{MgSO}_4$  to suppress  $I_{\text{Ca,L}}$  amplitude in both untreated and SGX-treated cells (data not shown). Previous observations have shown that DEX did not increase the activity of large-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  ( $\text{BK}_{\text{Ca}}$ ) channels in pituitary cells treated with methyl- $\beta$ -cyclodextrin [30],

suggesting that the binding of DEX to the protein(s) of  $BK_{Ca}$  channels relies on membrane cholesterol.

Whether DEX produces any significant effect on  $I_{Na}$  or  $I_{K(DR)}$  in NSC-34 cells or primary motor neurons needs to be further investigated. Whether the presence of SGX alters the cholesterol content in surface membranes and influences the DEX effect on  $I_{Ca,L}$  also remains to be explored. Nonetheless, our experimental results are consistent with earlier work showing that DEX is effective at exerting inhibitory effects on SGX reversal of rocuronium-induced neuromuscular block [28]. Alternatively, removal by SGX of DEX-induced block of  $I_{Ca,L}$  could be due mostly to the possibility that, similar to a mechanism by which it can reverse muscle relaxing effects by rocuronium, the SGX molecule can effectively encapsulate the DEX molecule [1, 3–5].

A recent report showed that methyl- $\beta$ -cyclodextrin, a cholesterol-depleting agent, could induce activation of matrix metalloproteinase-2 (MMP-2) [31]. However, the reduction by SGX treatment of  $I_{K(DR)}$  activation rate observed in differentiated NSC-34 cells was unable to be reversed by isobavachalcone (10  $\mu$ M) known to be an inhibitor of MMP-2 activity [32]. Therefore, alterations by SGX treatment of activation kinetics of  $I_{K(DR)}$  observed in differentiated NSC-34 cells are not closely associated with a mechanism linked to MMP-2 activation.

It should be noted that the pipette solution used in this study contained 3 mM ATP, which can adequately suppress the activity of ATP-sensitive  $K^+$  ( $K_{ATP}$ ) channels. The activity of  $K_{ATP}$  channels did not differ between the untreated and SGX-treated cells (data not shown). Changes by SGX treatment of  $I_{K(DR)}$  amplitude and gating observed in differentiated NSC-34 cells is unlikely to arise from inhibition of  $K_{ATP}$  channels.

The observed block of  $I_{K(DR)}$  caused by SGX treatment actually provides an intriguing mechanism for its inhibition that relies on the closed (resting) state of the  $K_v3.1$ -encoded channels. The  $K_v3.1$ -encoded currents were enriched in many central neurons including hippocampal pyramidal neurons, auditory neurons, and Purkinje cells [9, 11, 13, 14]. The activity of these  $K_v$  channels is recognized as participating in electrical behaviors of fast-spiking neurons [9, 10, 15, 16]. Challenging cells with SGX reduced the amplitude of  $I_{K(DR)}$  and slowed the activation time course of this current recorded from differentiated NSC-34 cells as well. The present observations would clearly initiate further studies to understand the SGX effects on electrical activity of motor neurons. Whether SGX-induced reversal of rocuronium-induced neuromuscular blockade is due partly to its blocking of  $I_{K(DR)}$  in motor neurons in vivo remains to be further investigated. Some adverse effect such as movement of a limb or the body may be partly explained by its inhibitory effect on  $I_{K(DR)}$ .

It is noted that neostigmine, an inhibitor of acetylcholinesterase activity, is a typical drug used in anesthesia for reversing the effect of rocuronium-induced neuromuscular blockade. In our study, neostigmine (1  $\mu$ M) did not exert any effect on the amplitude and gating of  $I_{K(DR)}$  in differentiated NSC-34 cells (Additional file 1: Fig. S1). Findings from our study might explain previous observations showing that SGX could reverse more rapidly rocuronium-induced neuromuscular blockade [33] or that the treatment with SGX was associated with less frequent dry mouth than that of neostigmine [34]. Therefore, it remains to be further delineated whether SGX might exert differential actions when it is used with patients who have been administered with DEX or other glucocorticoids [4], if similar findings presented here occur in vivo. Nonetheless, as motor neurons are exposed to SGX, the amplitude and gating of  $I_{K(DR)}$  could be modified and these actions might significantly contribute to functional activities of motor neurons.

## Conclusion

The SGX treatment may influence the amplitude and gating of  $I_{K(DR)}$  and its actions could contribute to functional activities of motor neurons if similar findings occurred in vivo.

## Additional files

**Additional file 1.** Lack of effect of neostigmine on  $I_{K(DR)}$  recorded from differentiated NSC-34 neuronal cells. In these experiments, cells were bathed in  $Ca^{2+}$ -free Tyrode's solution containing 1  $\mu$ M tetrodotoxin and 0.5 mM  $CdCl_2$ , and the recording pipette was filled with  $K^+$ -containing solution. (A) Original current trace obtained in the absence (blue) and presence (red) of 1  $\mu$ M neostigmine. Inset indicates the voltage protocol used. (B) Bar graph showing no significant effect of neostigmine (1  $\mu$ M) on  $I_{K(DR)}$  amplitude measured at the end of depolarizing pulses (mean  $\pm$  SEM;  $n = 8$  for each bar;  $P > 0.05$ ).

**Additional file 2.** Time course of  $I_{K(DR)}$  elicited by 100-ms depolarizing pulse from  $-50$  to  $+50$  mV (A) or repetitive cumulative activation (B) which was taken from the same NSC-34 cell. The number of raw data (indicated in filled circles) was reduced for clarity. Inset in (A) indicates the voltage protocol used. In (B), each  $I_{K(DR)}$  was evoked by 10-ms repetitive pulses to  $+50$  mV, each of which lasted 10 ms with 5-ms interval. The activation time constants taken from (A) and (B) (indicated in smooth gray lines) are 25.4 and 19.5 ms, respectively.

**Additional file 3.** Effect of nifedipine and  $\omega$ -conotoxin GVIA on  $I_{Ca,L}$  in differentiated NSC-34 neuronal cells. In these experiments, cells were bathed in normal Tyrode's solution containing 1  $\mu$ M tetrodotoxin and the recording pipette was filled with  $Cs^+$ -containing solution. (A) Original  $I_{Ca,L}$  trace obtained in the absence (blue) and presence (red) of 1  $\mu$ M nifedipine. Inset indicates the voltage protocol used. (B) Bar graph showing effect of nifedipine (1  $\mu$ M) and  $\omega$ -conotoxin GVIA (1  $\mu$ M) on the peak amplitude of  $I_{Ca,L}$  in these cells ( $n = 7$ ; mean  $\pm$  SEM for each bar). \* Significantly different from control ( $P < 0.05$ ).

## Abbreviations

SGX: sugammadex;  $I_{K(DR)}$ : delayed-rectifier  $K^+$  current;  $I_{Na}$ : voltage-gated  $Na^+$  current;  $I_{Ca,L}$ : L-type  $Ca^{2+}$  current; DEX: dexamethasone; TEA: tetraethylammonium chloride; FBS: fetal bovine serum; DMEM: Dulbecco's modified Eagle medium;  $I-V$ : current versus voltage;  $\Delta G_0$ : the free energy involved in the

gating of  $I_{K(DR)}$ ; SEM: standard error of the mean;  $\Delta\Delta G_0$ : the perturbation by SGX treatment of free energy;  $BK_{Ca}$ : large-conductance  $Ca^{2+}$ -activated  $K^+$ ; MMP-2: matrix metalloproteinase-2;  $K_{ATP}$  channel: ATP-sensitive  $K^+$  channel.

#### Authors' contributions

H-TH performed the research and analyzed the data; H-TH, Y-CL and S-NW designed the research study and wrote the paper; Yi-CL, Y-MH, Y-TT, and S-NW contributed essential reagents and tools. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup> Graduate Institute of Medicine, Kaohsiung Medical University, Kaohsiung City 80708, Taiwan. <sup>2</sup> Department of Anesthesia, Kaohsiung Municipal Ta-Tung Hospital, Kaohsiung City 80145, Taiwan. <sup>3</sup> Department of Pharmacology, School of Medicine, Kaohsiung Medical University, Kaohsiung City 80708, Taiwan. <sup>4</sup> Department of Physiology, National Cheng Kung University Medical College, No. 1, University Road, Tainan City 70101, Taiwan. <sup>5</sup> Graduate Institute of Natural Products, School of Pharmacy, Kaohsiung Medical University, Kaohsiung City 80708, Taiwan.

#### Acknowledgements

We thank Ming-Chun Hsu, Yan-Ming Huang and Huei-Jen Chen for their assistance in cell preparations. We also appreciated the English editing by Steve Tredrea.

#### Competing interests

The authors declare that they have no competing interests.

#### Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

#### Funding

Funding was provided by Kaohsiung Medical University (Award Number: kmth-104-051).

Received: 22 March 2016 Accepted: 8 December 2016

Published online: 03 January 2017

#### References

- Bom A, Hope F, Rutherford S, Thomson K. Preclinical pharmacology of sugammadex. *J Crit Care*. 2009;24(1):29–35.
- Paton L, Gupta S, Blacoe D. Successful use of sugammadex in a 'can't ventilate' scenario. *Anaesthesia*. 2013;68(8):861–4.
- Hudson ME, Rietbergen H, Chelly JE. Sugammadex is effective in reversing rocuronium in the presence of antibiotics. *BMC Anesthesiol*. 2014;14:69.
- Ledowski T. Sugammadex: what do we know and what do we still need to know? A review of the recent (2013 to 2014) literature. *Anaesth Intensive Care*. 2015;43(1):14–22.
- Brueckmann B, Sasaki N, Grobara P, Li M, Woo T, de Bie J, Maktabi M, Lee J, Kwo J, Pino R, Sabouri AS, McGovern F, Staehr-Rye AK, Eikermann M. Effects of sugammadex on incidence of postoperative residual neuromuscular blockade: a randomized, controlled study. *Br J Anaesth*. 2015;115(5):743–51.
- Palanca JM, Aguirre-Rueda D, Granell MV, Aldasoro M, Garcia A, Iradi A, Obrador E, Mauricio MD, Vila J, Gil-Bisquert A, Valles SL. Sugammadex, a neuromuscular blockade reversal agent, causes neuronal apoptosis in primary cultures. *Int J Med Sci*. 2013;10(10):1278–85.
- Wachi M, Uehara K, Fujinaka W, Takatori M, Tada K. Use of sugammadex in a patient with amyotrophic lateral sclerosis. *Masui*. 2011;60(12):1408–10.
- Takeuchi R, Hoshijima H, Doi K, Nagasaka H. The use of sugammadex in a patient with Kennedy's disease under general anesthesia. *J Anaesth*. 2014;8(3):418–20.
- Rudy B, McBain CJ. Kv3 channels: voltage-gated  $K^+$  channels designed for high-frequency repetitive firing. *Trends Neurosci*. 2001;24(9):517–26.
- Baranauskas G. Ionic channel function in action potential generation: current perspective. *Mol Neurobiol*. 2007;35(2):129–50.
- Lien CC, Jonas P. Kv3 potassium conductance is necessary and kinetically optimized for high-frequency action potential generation in hippocampal interneurons. *J Neurosci*. 2003;23(6):2058–68.
- Lin MW, Wang YJ, Liu SJ, Lin AA, Lo YC, Wu SN. Characterization of aconitine-induced block of delayed rectifier  $K^+$  current in differentiated NG108-15 neuronal cells. *Neuropharmacology*. 2008;54(6):912–23.
- Puente N, Mendizabal-Zubiaga J, Elezgarai I, Reguero L, Buceta I, Grandes P. Precise localization of the voltage-gated potassium channel subunits Kv3.1b and Kv3.3 revealed in the molecular layer of the rat cerebellar cortex by a pre-embedding immunogold method. *Histochem Cell Biol*. 2010;134(4):403–9.
- Strumbos JG, Polley DB, Kaczmarek LK. Specific and rapid effects of acoustic stimulation on the tonotopic distribution of Kv3.1b potassium channels in the adult rat. *Neuroscience*. 2010;167(3):567–72.
- Ding S, Matta SG, Zhou FM. Kv3-like potassium channels are required for sustained high-frequency firing in basal ganglia output neurons. *J Neurophysiol*. 2011;105(2):554–70.
- Hoppa MB, Gouzer G, Armbruster M, Ryan TA. Control and plasticity of the presynaptic action potential waveform at small CNS nerve terminals. *Neuron*. 2014;84(4):778–89.
- Muona M, Berkovic SF, Dibbens LM, Oliver KL, Maljevic S, Bayly MA, Joensuu T, Canafoglia L, Franceschetti S, Michelucci R, Heron SE, Hildebrand MS, Andermann E, Andermann F, Gambardella A, Tinuper P, Licchetta L, Scheffer IE, Criscuolo C, Filla A, Feriazzo E, Ahmad J, Ahmad A, Baykan B, Said E, Topcu M, Riguzzi P, King MD, Ozkara C, Andrade DM, Engelsen BA, Crespel A, Lindenau M, Lohmann E, Saletti V, Massano J, Privitera M, Espay AJ, Kauffmann B, Duchowny M, Meller RS, Straussberg R, Afawi Z, Ben-Zeev B, Samocha KE, Daly MJ, Petrou S, Lerche H, Palotie A, Lehesjoki AW. A recurrent de novo mutation in KCNC1 causes progressive myoclonus epilepsy. *Nat Genet*. 2015;47(1):39–46.
- Cashman NR, Durham HD, Blusztajn JK, Oda K, Tabira T, Shaw LT, Dahrouge S, Antel JP. Neuroblastoma × spinal cord (NSC) hybrid cell lines resemble developing motor neurons. *Dev Dyn*. 1992;194(3):209–21.
- Hsu HT, Tseng YT, Lo YC, Wu SN. Ability of naringenin, a bioflavonoid, to activate M-type potassium current in motor neuron-like cells and to increase  $BK_{Ca}$ -channel activity in HEK293T cells transfected with  $\alpha$ -hSlo subunit. *BMC Neurosci*. 2014;15:135.
- Eggett CJ, Crosier S, Manning P, Cookson MR, Menzies FM, McNeil CJ, Shaw PJ. Development and characterization of a glutamate-sensitive motor neuron cell line. *J Neurochem*. 2000;74(5):1895–902.
- So EC, Wu KC, Kao FC, Wu SN. Effects of midazolam on ion currents and membrane potential in differentiated motor neuron-like NSC-34 and NG108-15 cells. *Eur J Pharmacol*. 2014;724:152–60.
- Wu SN, Hsu MC, Liao YK, Wu FT, Jong YJ, Lo YC. Evidence for inhibitory effects of flupirtine, a centrally acting analgesic, on delayed rectifier  $K^+$  currents in motor neuron-like cells. *Evid Based Complement Alternat Med*. 2012;2012:148403.
- Huang CW, Wu YJ, Wu SN. Modification of activation kinetics of delayed rectifier  $K^+$  currents and neuronal excitability by methyl- $\beta$ -cyclodextrin. *Neuroscience*. 2011;176:431–41.
- Yifrach O, MacKinnon R. Energetics of pore opening in a voltage-gated  $K^+$  channel. *Cell*. 2002;111(2):231–9.
- Wu SN, Yeh CC, Huang HC, So EC, Lo YC. Electrophysiological characterization of sodium-activated potassium channels in NG108-15 and NSC-34 motor neuron-like cells. *Acta Physiol*. 2012;206(2):120–34.
- Wu SN, Yeh CC, Huang HC, Yang WH. Cholesterol depletion with (2-hydroxypropyl)- $\beta$ -cyclodextrin modifies the gating of membrane electroporation-induced inward current in pituitary tumor GH<sub>3</sub> cells: experimental and analytical studies. *Cell Physiol Biochem*. 2011;28(5):959–68.
- Chen BS, Lo YC, Peng H, Hsu TI, Wu SN. Effects of ranolazine, a novel anti-anginal drug, on ion currents and membrane potential in pituitary tumor GH<sub>3</sub> cells and NG108-15 neuronal cells. *J Pharmacol Sci*. 2009;110(3):295–305.
- Rezonja K, Sostaric M, Vidmar G, Mars T. Dexamethasone produces dose-dependent inhibition of sugammadex reversal in in vitro innervated primary human muscle cells. *Anesth Analg*. 2014;118(4):755–63.
- Huang MH, So EC, Liu YC, Wu SN. Glucocorticoids stimulate the activity of large-conductance  $Ca^{2+}$ -activated  $K^+$  channels in pituitary GH<sub>3</sub> and AtT-20 cells via a non-genomic mechanism. *Steroids*. 2006;71(2):129–40.

30. Lin MW, Wu AZ, Ting WH, Li CL, Cheng KS, Wu SN. Changes in membrane cholesterol of pituitary tumor (GH<sub>3</sub>) cells regulate the activity of large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channels. *Chin J Physiol*. 2006;49(1):1–13.
31. Kim S, Oh JH, Lee Y, Lee J, Cho KH, Chung JH. Induction of tissue inhibitor of matrix metalloproteinase-2 by cholesterol depletion leads to the conversion of proMMP-2 into active MMP-2 in human dermal fibroblasts. *Exp Mol Med*. 2010;42(1):38–46.
32. Ngameni B, Touaibia M, Belkaid A, Ambassa P, Watchueng J, Patnam R, Ngadjui BT, Annabi B, Roy R. Inhibition of matrix metalloproteinase-2 secretion by chalcones from the twigs of *Dorstenia barteri* Bureau. *ARKIVOC*. 2007;9:91–103.
33. Abrishami A, Ho J, Wong J, Yin L, Chung F. Cochrane corner: sugammadex, a selective reversal medication for preventing postoperative residual neuromuscular blockade. *Anesth Analg*. 2010;110(4):1239.
34. Sacan O, White PF, Tufanogullari B, Klein K. Sugammadex reversal of rocuronium-induced neuromuscular blockade: a comparison with neostigmine-glycopyrrolate and edrophonium-atropine. *Anesth Analg*. 2007;104(3):569–74.

Submit your next manuscript to BioMed Central  
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

